

Ministry of Basic and Senior Secondary Education

Pupils' Handbook for Senior Secondary Mathematics



DOI: 10.5281/zenodo.3745342 Please see final page for further information.

Foreword

These Lesson Plans and the accompanying Pupils' Handbooks are essential educational resources for the promotion of quality education in senior secondary schools in Sierra Leone. As Minister of Basic and Senior Secondary Education, I am pleased with the professional competencies demonstrated by the writers of these educational materials in English Language and Mathematics.

The Lesson Plans give teachers the support they need to cover each element of the national curriculum, as well as prepare pupils for the West African Examinations Council's (WAEC) examinations. The practice activities in the Pupils' Handbooks are designed to support self-study by pupils, and to give them additional opportunities to learn independently. In total, we have produced 516 lesson plans and 516 practice activities – one for each lesson, in each term, in each year, for each class. The production of these materials in a matter of months is a remarkable achievement.

These plans have been written by experienced Sierra Leoneans together with international educators. They have been reviewed by officials of my Ministry to ensure that they meet the specific needs of the Sierra Leonean population. They provide step-by-step guidance for each learning outcome, using a range of recognized techniques to deliver the best teaching.

I call on all teachers and heads of schools across the country to make the best use of these materials. We are supporting our teachers through a detailed training programme designed specifically for these new lesson plans. It is really important that the Lesson Plans and Pupils' Handbooks are used, together with any other materials they may have.

This is just the start of educational transformation in Sierra Leone as pronounced by His Excellency, the President of the Republic of Sierra Leone, Brigadier Rtd Julius Maada Bio. I am committed to continue to strive for the changes that will make our country stronger and better.

I do thank our partners for their continued support. Finally, I also thank the teachers of our country for their hard work in securing our future

Mr. Alpha Osman Timbo

Minister of Basic and Senior Secondary Education

The policy of the Ministry of Basic and Senior Secondary Education, Sierra Leone, on textbooks stipulates that every printed book should have a lifespan of three years.

To achieve thus, DO NOT WRITE IN THE BOOKS.

Table of Contents

Lesson 49: Powers and Roots of Logarithms – Numbers greater than 1	3
Lesson 50: Logarithms – Numbers less than 1	6
Lesson 51: Antilogarithms – Numbers less than 0	8
Lesson 52: Multiplication and Division using Logarithms – Numbers less than 1	10
Lesson 53: Powers and Roots of Logarithms – Numbers less than 1	12
Lesson 54: Laws of Logarithms – Part 1	15
Lesson 55: Laws of Logarithms – Part 2	18
Lesson 56: Laws of Logarithms – Part 3	20
Lesson 57: Define and Describe Sets and Elements of a Set	23
Lesson 58: Define and Describe Sets and Elements of a Set	26
Lesson 59: Finite and Infinite Sets	28
Lesson 60: Null/Empty, Unit and Universal Sets	30
Lesson 61: Equivalent and Equal Sets	32
Lesson 62: Subsets	34
Lesson 63: Intersection of 2 Sets	37
Lesson 64: Intersection of 3 Sets	40
Lesson 65: Disjoint Sets	43
Lesson 66: Union of Two Sets	45
Lesson 67: Complement of a Set	48
Lesson 68: Problem Solving with 2 Sets	51
Lesson 69: Problem Solving with 3 Sets – Part 1	55
Lesson 70: Problem Solving with 3 Sets – Part 2	58
Lesson 71: Use of Variables	61
Lesson 72: Simplification – Grouping Terms	64
Lesson 73: Simplification – Removing Brackets	67
Lesson 74: Simplification – Expanding Brackets	69
Lesson 75: Factoring – Common Factors	72
Lesson 76: Factoring – Grouping	74

Lesson 77: Substitution of Values	76
Lesson 78: Addition of Algebraic Fractions	78
Lesson 79: Subtraction of Algebraic Fractions	81
Lesson 80: Linear Equations	84
Lesson 81: Linear Equations with Brackets	87
Lesson 82: Linear Equations with Fractions	90
Lesson 83: Word Problems	93
Lesson 84: Substitution in Formulae	96
Lesson 85: Change of Subject – Part 1	99
Lesson 86: Change of Subject – Part 2	102
Lesson 87: Reduction to Basic Form of Surds	105
Lesson 88: Addition and Subtraction of Surds – Part 1	108
Lesson 89: Addition and Subtraction of Surds – Part 2	110
Lesson 90: Properties of Surds	112
Lesson 91: Multiplication of Surds – Part 1	114
Lesson 92: Multiplication of Surds – Part 2	117
Lesson 93: Rationalisation of the Denominator of Surds – Part 1	120
Lesson 94: Rationalisation of the Denominator of Surds – Part 2	122
Lesson 95: Expansion and Simplification of Surds	125
Lesson 96: Practice of Surds	127
Answer Key: Term 2	130
Appendix I: Logarithm Table	147
Appendix II: Anti-Logarithm Table	148

Introduction to the Pupils' Handbook

1

2

3

5

6

7

8

These practice activities are aligned to the Lesson Plans, and are based on the National Curriculum and the West Africa Examination Council syllabus guidelines. They meet the requirements established by the Ministry of Basic and Senior Secondary Education.



Learning Outcomes

1

Do not worry if you do not get the right answer -

ask for help and continue practising!

KEY TAKEAWAYS FROM SIERRA LEONE'S PERFORMANCE IN WEST AFRICAN SENIOR SCHOOL CERTIFICATE EXAMINATION – GENERAL MATHEMATICS¹

This section, seeks to outline key takeaways from assessing Sierra Leonean pupils' responses on the West African Senior School Certificate Examination. The common errors pupils make are highlighted below with the intention of giving teachers an insight into areas to focus on, to improve pupil performance on the examination. Suggestions are provided for addressing these issues.

Common errors

- 1. Errors in applying principles of BODMAS
- 2. Mistakes in simplifying fractions
- 3. Errors in application of Maths learned in class to real-life situations, and vis-aversa.
- 4. Errors in solving geometric constructions.
- 5. Mistakes in solving problems on circle theorems.
- 6. Proofs are often left out from solutions, derivations are often missing from quadratic equations.

Suggested solutions

- 1. Practice answering questions to the detail requested
- 2. Practice re-reading questions to make sure all the components are answered.
- 3. If possible, procure as many geometry sets to practice geometry construction.
- 4. Check that depth and level of the lesson taught is appropriate for the grade level.

¹ This information is derived from an evaluation of WAEC Examiner Reports, as well as input from WAEC examiners and Sierra Leonean teachers.

Lesson Title: Powers and roots of logarithms – Numbers greater than 1	Theme: Numbers and numeration	
Practice Activity: PHM1-L049	Class: SSS 1	

By the end of the lesson, you will be able to calculate powers and roots of numbers greater than 1 using logarithms.

Overview

This lesson is on calculating powers and roots of numbers greater than 1 using logarithms. For example, consider the problems 138^2 and $\sqrt[3]{46.5}$. These seem difficult to calculate without using a calculator. However, we can calculate them using logarithms.

The product rule is applied to evaluate powers of numbers greater than 1:

- Step 1. Find the logarithm of the number.
- Step 2. Multiply the logarithm by the power.
- Step 3. Find the antilog of the result in step 2.

Similarly, the **quotient rule** is applied to evaluate **roots** of numbers greater than 1:

- Step 1. Find the logarithm of the number.
- Step 2. Divide the logarithm by the root.
- Step 3. Find the antilogarithm of the result in step 2.

You can apply logarithms to solve problems with more than one operation. For example, $245.5^2 \times 1.08^3$. This involves powers and multiplication. We apply the order of operations (BODMAS). Find the logarithms of the numbers first, then apply all of the operations necessary before applying the antilogarithm to the result.

Solved Examples

1. Evaluate 138²

Solution:

Steps	Numbers	Logarithm
1. Find the logarithm	138	2.1399
2. Multiply by the power	138 ²	$2.1399 \times 2 = 4.2798$
3. Find the antilog		antilog 4.2798 = 19040
Answer	$138^2 = 19040$	

2. Evaluate 3.12⁵

Solution:

Steps	Numbers	Logarithm
1.	3.12	0.4942
2.	3.12 ⁵	$0.4942 \times 5 = 2.471$
3.	antilog 2.471=295.8	
Answer	$3.12^5 = 295.8$	

3. Evaluate $\sqrt{5}$ **Solution:** is a square root, so the

This

root that we will divide by is 2.

Steps	Numbers	Logarithm
1. Find the logarithm	5	0.6990
2. Divide by the root	$\sqrt{5}$	$0.6990 \div 2 = 0.3495$
3. Find the antilog		antilog 0.3495=2.237
Answer	$\sqrt{5} = 2.237$,

4. Evaluate $\sqrt[3]{64.5}$

Solution:

This is a cubed root, so the root that we will divide by is 3.

Steps	Numbers	Logarithm
1.	64.5	1.8096
2.	³ √64.5	$\frac{1.8096}{3} = 0.6032$
3.		antilog 0.6032= 4.011
Answer	$\sqrt[3]{64.5} = 4.011$	

5. Evaluate $\sqrt[4]{(25.1)^3}$

Solution:

This problem has both a square and a root on the number. Apply the square first using multiplication. Then, apply the root using division.

Steps	Number	Logarithm
Find the logarithm of $(25.1)^3$	25.1	1.3997
	$(25.1)^3$	$1.3997 \times 3 = 4.1991$
Find $\sqrt[4]{(25.1)^3}$	$\sqrt[4]{(25.1)^3}$	$4.1991 \div 4 = 1.0498$
	Answer	Antilog 1.0498 = 11.21

6. Evaluate $\sqrt{245.5} \times 8.01^2$

Solution:

In this case, roots and powers appear together. Find the logarithm of each separately. Then, add the logarithms to find the product of $\sqrt{245.5}$ and 8.01^2 .

Steps	Number	Logarithm
Find the logarithm of $\sqrt{245.5}$	245.5	2.3901
	$\sqrt{245.5}$	$2.3901 \div 2 = 1.1951$
Find the logarithm of 8.01 ²	8.01	0.9036
	8.01 ²	$0.9036 \times 2 = 1.8072$
Multiply the 2 terms	$\sqrt{245.5} \times 8.01^2$	
Add their logarithms		1.1951
		+1.8072
		3.0023
	Answer	Antilog 3.0023 = 1,006

Therefore, we have $\sqrt{245.5} \times 8.01^2 = 1,006$.

Practice

Use logarithm tables to:

- 1. Evaluate 31.1²
- 2. Evaluate 3.39⁴
- 3. Evaluate $\sqrt{225}$
- 4. Evaluate $\sqrt[4]{(62.5)^2}$
- 5. Evaluate $2.45^3 \times \sqrt[5]{611}$
- 6. Evaluate $\sqrt[3]{999} \div 2.33^2$

Lesson Title: Logarithms – Numbers less than 1	Theme: Numbers and numeration
Practice Activity: PHM1-L050	Class: SSS 1

By the end of the lesson, you will be able to find the logarithms of numbers less than 1 using logarithm tables.

Overview

This lesson is on finding the logarithm of numbers less than 1 using logarithm tables. In other words, we are going to find the logarithm of decimal numbers which lie in the range 0 to 1.

An easy rule that works for decimals is that the characteristic is equal to the number of zeros trailing the decimal plus 1. Consider an example, 0.0314. The number has 1 zero after the decimal, so the characteristic is -2. The characteristic can also be found by writing the decimal number in standard form. Note that $0.0314 = 3.14 \times 10^{-2}$. The power of 10 gives the characteristic. When a decimal number should be expressed in its standard form $a \times 10^{-n}$, the characteristic of the logarithm is \bar{n} .

The characteristic is not written with a negative sign. We put the minus sign on top of the characteristic. Thus, we have $\overline{2}$, which is read as "**bar**" **2**, but not "minus" 2.

The mantissa is found using a logarithm table, in the same way as with number greater than 1. Write the mantissa after the characteristic.

Solved Examples

1. Write the following numbers in standard form and derive the characteristics of their logarithms from the standard form.

a. 0.121 b. 0.00212 c. 0.000561 d. 0.00410

Solutions:

	Number	Standard form	Characteristics of logarithm
a.	0.121	1.21×10^{-1}	1
b.	0.00212	2.12×10^{-3}	3
C.	0.000561	5.61×10^{-4}	4
d.	0.00410	4.10×10^{-3}	3

2. Using the simple rule that the number of zeros trailing the decimal point plus 1 gives the characteristic, find the characteristic of the logarithm of each of the following numbers:

a. 0.315 b. 0.00456 c. 0.000128 d. 0.0000100

Solutions:

	Number	Logarithm	Characteristics of logarithm
a.	0.315	log 0.315	$-(0+1)=\overline{1}$
b.	0.00456	log 0.00456	$-(2+1) = \overline{3}$
C.	0.000128	log 0.000128	$-(3+1) = \overline{4}$
d.	0.0000100	log 0.0000100	$-(4+1) = \overline{5}$

3. Find the logarithms of the following numbers using the logarithm table: a. 0.153 b. 0.00654 c. 0.000812 d. 0.0000210

Solutions:

	Number	Logarithm	
a.	0.153	log 0.153	= 1.1847
b.	0.00654	log 0.00654	= 3.8156
C.	0.000812	log 0.000812	= 4.9096
d.	0.0000210	log 0.0000210	$0 = \overline{5}.3222$

Practice

1. Write the following numbers in standard form. Then, write the characteristic of each.

a. 0.4525 b. 0.0607 c. 0.00498 d. 0.000989

2. Using the simple rule that the number of zeros trailing the decimal point plus 1 gives the characteristic, find the characteristic of the logarithm of each of the following numbers:

a. 0.135 b. 0.00546 c. 0.0786 d. 0.0000341

Find the logarithms of the following numbers:
 a. 0.123 b. 0.0987 c. 0.00654 d. 0.000899

Lesson Title: Antilogarithm – Numbers less than 0	Theme: Numbers and Numeration
Practice Activity: PHM1-L051	Class: SSS 1

By the end of the lesson, you will be able to find the antilogarithm of numbers less than 0 using antilogarithm tables.

Overview

Today's lesson is going to be on how to find the antilogarithms of the logarithms of numbers less than 0.

Antilogarithm is the exact opposite of the logarithm of a number. It brings the number back to its ordinary form from the logarithm form. So if $X = \log b$, then antilog X = b

To find the antilogarithms of numbers we use the antilogarithm tables, which give the inverse value of the mantissa to four figures. The characteristic is then used to position the decimal point in the value read from the mantissa. Write the number of zeros after the decimal based on the characteristic. Remember that there are 1 fewer zeros after the decimal than the value of the characteristic.

Solved Examples

1. Read the antilogarithms of the following logarithms of numbers.

```
a. \overline{1}. 9178 b. \overline{2}. 0451 c. \overline{6}. 3251 d. \overline{3}. 8547 e. \overline{3}. 2045 f. \overline{1}. 3749
```

Solutions:

	Antilog	Value read from using	Number of zeros	Answer
		mantissa antilog table	after the decimal	
a.	1.9178	8,275	0 zeros	0.8275
b.	2.0451	1,109	1 zero	0.01109
C.	6.3251	2,113	5 zeros	0.000002113
d.	3.8547	7,157	2 zeros	0.007157
e.	3.2045	1,602	2 zeros	0.001602
f.	1.3749	2,371	0 zeros	0.2371

Practice

Find the antilogarithm of the following logarithms of numbers:

- 1. 2.3067
- 2. 4.0101
- **3**. <u>1</u>.2011

4. 4.4145
 5. 6.1101
 6. 5.9999
 7. 2.6303
 8. 3.3344

Lesson Title: Multiplication and division	Theme: Numbers and Numeration
using logarithms – Numbers less than 1	
Practice Activity: PHM1-052	Class: SSS 1

By the end of the lesson, you will be able to multiply and divide numbers less than 1 using logarithms.

Overview

To find the **product** of numbers less than 1 using logarithm tables, follow these steps:

- Find the logarithms of the numbers
- Add the logarithms
- Find the antilog of the numbers resulting from the sum.

Similarly, to find the **quotient** of numbers less than 1 using logarithm tables, follow these steps:

- Find the logarithms of the numbers
- Subtract the logarithms
- Find the antilogarithm of the number resulting from the subtraction.

Solved Examples

1. Evaluate 0.056×0.95

Solution:

Number	Logarithm	
0.056	2.7482	
0.95	$+\bar{1}.9777$	
Add logs for	2.7259	
multiplication		
Antilog $\bar{2}$. 7259 = 0.05320		

2. Evaluate $\frac{0.0978}{0.06}$

Solution:

Number	Logarithm	
0.0978	2.9903	
0.06	$-\overline{2}.7782$	
Subtract logs	0.2121	
for division		
Antilog of $0.2121 = 1.63$		

3. Evaluate $\frac{0.0067 \times 0.037}{0.57}$

Solution:

Apply the normal order of operations. Multiply the numbers in the numerator (using addition of logs) before dividing by the denominator (using subtraction of logs).

Number	Logarithm
0.0067	3.8261
0.037	$+\bar{2}.5682$
Add logs for multiplication	4 .3943
0.57	$-\bar{1}.7559$
Subtract logs for division	4 .6384
Antilog $\overline{4}$. 6384 = 0.0004349	

Practice

Evaluate the following using logarithmic tables:

- 1. $\frac{0.01875}{0.0015}$
- 2. 0.099×0.005
- **3**. 0.491 × 0.23
- 4. $\frac{0.456 \times 0.56}{0.456 \times 0.56}$
- 2.5 2.5
- 5. 0.45×0.613
- 0.9981 6.
- 0.321

Lesson Title: Powers and roots of logarithms – Numbers less than 1	Theme: Numbers and Numeration
Practice Activity: PHM1-L053	Class: SSS 1

By the end of the lesson, you will be able to calculate powers and roots of numbers less than 1 using logarithms.

Overview

To evaluate the **powers** of numbers less than 1 using logarithm table, follow these steps:

- Find the logarithm of the number
- Multiply the logarithm found by the power
- Use the antilog table to find the antilog of the result to get your answer

To evaluate the **roots** of numbers less than 1 using logarithm tables, follow these steps:

- Find the logarithm of the number
- Divide the logarithm found by the root
- Use antilog table to find the antilog of the result to get your answer

Solved Examples

- 1. Evaluate: a. 0.0762⁴; b. 0.063² Solutions:
 - a. 0.0762⁴

Number	Logarithm	
0.0762	2.8820	
Multiply log by power	<u>× 4</u>	
	5.5280	
Antilog $\overline{5}.5280 = 0.00003373$		

Answer: $0.0762^4 = 0.00003373$

b. 0.063²

Number	Logarithm
0.063	2.7993
Multiply log by power	<u>× 2</u>
	3.5986
Antilog $\bar{3}$. 5986 = 0.003968	

Answer: $0.063^2 = 0.003968$

2. Evaluate: a. $\sqrt{0.0025}$; b. $\sqrt[3]{0.000216}$

Solutions:

a. √0.0025

Number	Logarithm
0.0025	3.979
For square root divide log by 2	$\frac{\overline{3.979}}{2} = \frac{\overline{4}}{2} + \frac{1.3979}{2}$ $= \overline{2}.6999$
Antilog $\overline{2}$. 6999 = 0.0501	

Answer: $\sqrt{0.0025} = 0.0501$

b. $\sqrt[3]{0.000216}$

Number	Logarithm	
0.000216	4 .3341	
For cube root divide log by 3	$\frac{\overline{4.3341}}{3} = \frac{\overline{6}}{3} + \frac{2.3341}{3} = \overline{2.7780}$	
Antilog $\overline{2}$. 7780 = 0.05998		

Answer: $\sqrt[3]{0.000216} = 0.05998$

- 3. Evaluate: a. $0.00125^{\frac{1}{3}}$; b. $0.0081^{\frac{2}{3}}$ Solutions:
 - a. Note that $0.00125^{\frac{1}{3}} = \sqrt[3]{0.00125}$. Thus, divide the logarithm by 3:

Number	Logarithm
0.00125	3.0969
	<u>÷ 3</u>
	ī.0323
Antilog $\bar{1}.0323 = 0.1$	077

Answer: $0.00125^{\frac{1}{3}} = 0.1077$

b. Note that $0.0081^{\frac{2}{3}} = \sqrt[3]{0.0081^2}$. Thus, multiply the logarithm by 2, then divide the result by 3.

Number	Logarithm
0.0081	3.9085
Multiply	<u>× 2</u>
	5.8170
Divide	÷ 3
	$\overline{2}.6057$
Antilog $\bar{2}.6057 = 0.0$	403

Answer: $0.0081^{\frac{2}{3}} = 0.0403$

Practice

- 1. Evaluate: a. 0.0459^2 ; b. 0.00125^3
- 2. Evaluate: a. $\sqrt[3]{0.625}$; b. $\sqrt{0.0109}$
- 3. Evaluate: a. $\frac{0.451^2}{\sqrt{0.516}}$; b. $\sqrt[4]{0.064} \times 0.27^2$
- 4. Evaluate: a. $0.0567^{\frac{1}{2}}$; b. $0.123^{\frac{4}{5}}$

Lesson Title: Laws of logarithms – Part 1	Theme: Numbers and Numeration
Practice Activity: PHM1-L054	Class: SSS 1

By the end of the lesson, you will be able to identify that $\log_{10}(pq) = \log_{10} p + \log_{10} q$.

Overview

This lesson is on the multiplication law of logarithms. Remember that with indices, when you multiply powers of the same base, you keep the base and add the exponent. For example: $y^2 \times y^3 = y^{2+3} = y^5$.

Note that logarithms are exponents, and when you multiply, you are going to add the logarithms. This means, the log of a product is the sum of the logs. For example, $\log_a xy = \log_a x + \log_a y$.

Note also that the log of a sum is NOT the sum of the logs. For example, $\log_a(x + y) \neq \log_a x + \log_a y$. The log of a sum cannot be simplified.

There are some other basic rules to keep in mind when simplifying logarithms. You may use these throughout the next few lessons:

- The logarithm of any number to the base of the same number is 1, that is $\log_a a = 1$.
- The logarithm of 1 to any base is 0, that is $\log_a 1 = 0$.

Solved Examples

1.	Simplify	the following:			
	a. log	$3 + \log 4$	b. log 9	+ log 2	c. $\log 3 + \log 2 + \log 5$
	Solut	ions:			
	a.	$\log 3 + \log 4$		$= \log(3 \times 4)$	4)
				$= \log 12$	
	b.	log 9 + log 2		$= \log(9 \times 2)$	2)
				$= \log 18$	
	C.	log 3 + log 2 +	log 5	$= \log(3 \times 2)$	2 × 5)
				$= \log 30$	
2.	Expand t	he following:			

- a. log₅ 18
- b. log₄ 14
- **c**. log₂ 35
- d. $\log_3(4m + 5)$

Solutions:

a. log ₅ 18	$= \log_5(6 \times 3)$ $= \log_5 6 + \log_5 3$
or: log ₅ 18	$= \log_5(9 \times 2)$ $= \log_5 9 + \log_5 2$
b. log ₄ 14	$= \log_4(2 \times 7)$ $= \log_4 2 + \log_4 7$
c . log ₂ 35	$= \log_2(5 \times 7)$ $= \log_2 5 + \log_2 7$

d. $\log_3(4m + 5)$: Cannot be expanded

3. Given that $\log_{10} 3 = 0.4771$, $\log_{10} 4 = 0.6021$, $\log_{10} 5 = 0.6990$ and $\log_{10} 6 = 0.7782$, find the values of the following:

a. log₁₀ 9 b. log₁₀ 24 c. log₁₀ 20 d. log₁₀ 30 e. log₁₀ 72 Solutions: b. $\log_{10} 24 = \log_{10} (6 \times 4)$ a. $\log_{10} 9$ $= \log_{10}(3 \times 3)$ $= \log_{10} 3 + \log_{10} 3$ $= \log_{10} 6 + \log_{10} 4$ = 0.4771 + 0.4771= 0.7782 + 0.6021= 0.9542= 1.3803c. $\log_{10} 20 = \log_{10} (4 \times 5)$ d. log₁₀ 30 $= \log_{10}(5 \times 6)$ $= \log_{10} 4 + \log_{10} 5$ $= \log_{10} 5 + \log_{10} 6$ = 0.6021 + 0.6990= 0.6990 + 0.7782= 1.4772= 1.3011 e. $\log_{10} 72 = \log_{10} (3 \times 4 \times 6)$

 $= \log_{10} 3 + \log_{10} 4 + \log_{10} 6$ = 0.4771 + 0.6021 + 0.7782= 1.8574

Practice

1.	Expand the following	:		
	a. log ₁₀ 14 b. lo	g ₁₀ 15	c . log 49	d. log 25
2.	Simplify:			
	a. $\log_2 6 + \log_2 6$	g ₂ 8		
	b. $\log_5 4 + \log_5 4$	$g_5 2 + l_{10}$	0g ₅ 3	
	c. $\log_4 8 + \log_4 8$	$g_4 8 + l$	og ₄ 2	
	d. $\log_3 9 + \log_3 9$	$g_3 2 + l_2$	og ₃ 2	

- 3. Given that $\log_{10} 2 = 0.3010$, $\log_{10} 5 = 0.6990$, and $\log_{10} 7 = 0.8451$, find the value of the following:
 - a. log 28
 - b. log 56
 - c. log 70

Lesson Title: Laws of logarithms – Part	Theme: Numbers and Numeration
2	
Practice Activity: PHM1-L055	Class: SSS 1

By the end of the lesson, you will be able to identify that $\log_{10}\left(\frac{p}{q}\right) = \log_{10} p - \log_{10} q$.

Overview

This lesson is on the division law of logarithms. Remember that with indices, when you divide powers of the same base, you keep the base and subtract the exponent. For example: $y^7 \div y^3 = y^{7-3} = y^4$.

Logarithms are also a way of expressing exponents. So when you take the logarithm of the quotient of any two numbers, it is equal to simply subtracting their logarithms of the same base. That is, $\log_a \left(\frac{x}{y}\right) = \log_a x - \log_a y$.

Note that the log of a difference is NOT the difference of the logs. That is, $\log_a(x - y) \neq \log_a x - \log_a y$. The log of a difference cannot be simplified.

Solved Examples

1. Expand: a. $\log_{10} 3\frac{1}{3}$ b. $\log_{10} 9\frac{5}{4}$ Solutions:

First, write each mixed fraction as an improper fraction of the form $\frac{x}{y}$. Then expand using the division law of logarithms.

a.
$$\log_{10} 3\frac{1}{3} = \log_{10} \left(\frac{10}{3}\right)$$

= $\log_{10} 10 - \log_{10} 3$
b. $\log_{10} 9\frac{4}{5} = \log_{10} \left(\frac{49}{5}\right)$
= $\log_{10} 49 - \log_{10} 5$

2. Simplify: a. $\log_3 21 - \log_3 7$ b. $\log_3 27 - \log_3 18$ Solutions:

a.
$$\log_3 21 - \log_3 7 = \log_3 \left(\frac{21}{7}\right)$$
 b. $\log_3 27 - \log_3 18 = \log_3 \left(\frac{27}{18}\right)$
= $\log_3 3$ = $\log_3 \left(\frac{3}{2}\right)$

3. Given that $\log_5 2 = 0.431$ and $\log_5 3 = 0.682$, find the value of a. $\log_5\left(1\frac{2}{3}\right)$ b. $\log_5\left(1\frac{1}{2}\right)$

Solutions:

First, rewrite each mixed fraction as an improper fraction. Then, expand and solve by substituting the given values.

- a. $\log_5\left(1\frac{2}{3}\right) = \log_5\left(\frac{5}{3}\right)$ $= \log_5 5 - \log_5 3$ = 1 - 0.682 = 0.318b. $\log_5\left(1\frac{1}{2}\right) = \log_5\left(\frac{3}{2}\right)$ $= \log_5 3 - \log_5 2$ = 0.682 - 0.431= 0.251
- 4. Given that $\log_{10} 7 = 0.8451$, $\log_{10} 2 = 0.3010$ and $\log_{10} 3 = 0.4771$, find the value of: a. $\log_{10} \left(\frac{9}{7}\right)$ b. $\log_{10} 0.21$ Solutions:
 - a. $\log_{10}\left(\frac{9}{7}\right)$ $= \log_{10}9 - \log_{10}7$ $= \log_{10}3 \times 3 - \log_{10}7$ $= \log_{10}3 + \log_{10}3 - \log_{10}7$ = 0.4771 + 0.4771 - 0.8451 = 0.1091b. $\log_{10}0.21 = \log\frac{21}{100}$ $= \log_{10}21 - \log_{10}100$ $= \log_{10}(7 \times 3) - \log_{10}(10 \times 10)$ $= \log_{10}7 + \log_{10}3 - (\log_{10}10 + \log_{10}10)$ = 0.8451 + 0.4771 - (1 + 1)= -0.6778

Practice

- 1. Expand: a. $\log_7 5\frac{2}{3}$ b. $\log_9 7\frac{4}{5}$
- 2. Simplify: a. $\log_4 32 \log_4 8$ b. $\log_3 48 \log_3 27$
- 3. Given that $\log_5 2 = 0.431$ and $\log_5 3 = 0.682$ find:

a.
$$\log_5 6\frac{2}{3}$$
 b. $\log_5 0.6$ c. $\log_5 \left(\frac{9}{25}\right)$

4. Given that $\log_{10} 7 = 0.8451$, $\log_{10} 2 = 0.3010$, $\log_{10} 3 = 0.4771$ find: a. $\log_{10} \left(11\frac{4}{6}\right)$ b. $\log_{10} 6\frac{1}{8}$

Lesson Title: Laws of logarithms – Part	Theme: Numbers and Numeration
3	
Practice Activity: PHM1-L056	Class: SSS 1

By the end of the lesson, you will be able to identify that $\log_{10}(P)^n = n \log_{10} P$.

Overview

When evaluating certain logarithmic expressions and equations, the logarithm power law can be used. The logarithm power/exponent law states that we can transfer the exponent of a number whose logarithm we are finding to the front of logarithm. The general rule is $\log_a m^r = r \log_a m$.

Along with the product and quotient rule, the logarithm power rule can be used for expanding and simplifying logarithmic expressions.

Note that if you have an entire logarithmic expression raised to an exponent, it cannot be simplified using this rule. In other words, $(\log_a m)^x \neq x \log_a m$.

Solved Examples

1. Simplify the following expressions:

a. $\log_{10} 2^7$ b. $\log_5 a^3$ c. $\log_8 8^4$ d. $\log_2 4$

Solutions:

Rewrite each expression using the third law of logarithms:

- a. $\log_{10} 2^7 = 7 \log_{10} 2$
- b. $\log_5 a^3 = 3 \log_5 a$
- c. $\log_8 8^4 = 4 \log_8 8 = 4 \times 1 = 4$
- d. For this example, first rewrite 4 as an index: 2^{2}

 $\log_2 4 = \log_2 2^2$ Now apply the third law of logarithms: $\log_2 2^2 = 2 \log_2 2 = 2 \times 1 = 2$

2. Given $\log_{10} 3 = 0.4771$ and $\log_{10} 2 = 0.3010$ evaluate: a. $\log_{10} 32$ b. $\log_{10} 27$ c. $\log_{10} 36$

Solutions:

a. $\log_{10} 32 = \log_{10} 2^5$ = $5 \log_{10} 2$ = 5(0.3010)= 1.505b. $\log_{10} 27 = \log_{10} 3^3$

=
$$3 \log_{10} 3$$

= $3(0.4771)$
= 1.4313
c. $\log_{10} 36 = \log_{10}(4 \times 9) = \log_{10} 4 + \log_{10} 9$
= $\log_{10} 2^2 + \log_{10} 3^2$
= $2 \log_{10} 2 + 2 \log_{10} 3$
= $2(0.3010) + 2(0.4771) = 1.5562$

3. Simplify:

a.
$$\frac{\log_a 8 + \log_a 16 - \log_a 2}{\log_a 32}$$

log₂ 8³

b.
$$\frac{\log_2 8^3}{\log_2 8}$$

Solutions:

a.
$$\frac{\log_a 8 + \log_a 16 - \log_a 2}{\log_a 32} = \frac{\log_a \left(\frac{8 \times 16}{2}\right)}{\log_a 32} = \frac{\log_a 64}{\log_a 32} = \frac{\log_a 2^6}{\log_a 2^5} = \frac{6(\log_a 2)}{5(\log_a 2)} = \frac{6}{5} = 1\frac{1}{5}$$

b.
$$\frac{\log_2 8^3}{\log_2 8} = \frac{3(\log_2 8)}{(\log_2 8)} = 3$$

- 4. Simplify:
 - a. log₁₀ 25 + log₁₀ 4
 b. log₃ 15 + log₃ 9 log₃ 5
 Solutions:
 - a.
- $log_{10} 25 + log_{10} 4 = log_{10} (25 \times 4)$ = log_{10} 100 = log_{10} 10² = 2 log_{10} 10 = 2 \times 1 = 2

b.

$$log_{3} 15 + log_{3} 9 - log_{3} 5 = log_{3} \left(\frac{15 \times 9}{5}\right)$$

= log_{3} 27
= log_{3} 3^{3}
= 3(log_{3} 3)
= 3 \times 1 = 3

Practice

- 1. Simplify: a. $\log_3 x^8$ b. $\log_2 8$ c. $\log_3 3^{17}$
- 2. Given $\log_5 2 = 0.431$ and $\log_5 3 = 0.682$ find the values of:
 - a. $\log_5 8$
 - b. $\log_5\left(\frac{4}{5}\right)$
 - c. log₅ 18

3. Simplify:

a.
$$\frac{\log_{10} 625}{\log_{10} 25}$$

b. $\frac{\log_{10} a^4 - \log_{10} a^2}{\log_{10} a^3}$
c. $\frac{\log_{10} 81}{\log_{10} 3}$

4. Evaluate: a. $3 + \log_3 81$ b. $(\log_5 125)^2 - \log_5 25$

Lesson Title: Define and describe sets	Theme: Numbers and Numeration
and elements of a set	
Practice Activity: PHM1-L057	Class: SSS 1

By the end of the lesson, you will be able to use various ways of writing and describing sets in terms of their numbers and elements.

Overview

A set is a collection of objects that have something in common or follow a common rule. For example, a set of fruits can be mangoes, oranges, guavas, pears, and so on.

The objects or things which make up a set are called **elements** of the set. In writing or defining sets, the symbol \in means "is a member of". For example, if we represent the set of even numbers between 0 and 10 by A, then we have A = {2, 4, 6, 8}. We can write $2\in A$ to show that 2 is a member of the set A.

The symbol \notin means "is not a member of" so $3\notin A$ means 3 is not a member of set A. We usually use a capital letter for the name of the set (A, B, C, ...) and sometimes use lowercase letters (a, b, c, ...) for the elements of the set.

The "cardinality of a set" or the "cardinal number of a set" is the number of elements in the set. It is denoted with the letter n and brackets. For example, consider n(A) = 4. This statement says that the set A has 4 elements.

Sometimes the members of a set may be too many for us to list them all. For example, the set *B* of integers from 1 to 1000. It is tedious to list all its members so we usually list few of the first elements and then continue with three dots (...) and write the last number. The set *B* can be written $B = \{1, 2, 3, 4, ..., 1000\}$

Solved Examples

- 1. Define each of the following sets of things by their common property
 - a. Pots, spoons, frying pan, cups, plates.
 - b. Paper clip, paper, stapler, pen, pencil, envelope.
 - c. 1, 4, 9, 16, 25, 36.

Solutions:

- a. Pots, spoons, frying pan, cups, plates are a set of eating utensils.
- b. Paper clip, paper, stapler, pen, pencil, envelope are a set of stationary or office materials.
- c. 1, 4, 9, 16, 25, 36 are a set of integers of perfect squares between 0 and 40.

- 2. List the elements of the following sets:
 - a. {The months of the years that have 30 days}
 - b. {All the even numbers between 0 and 100}
 - c. {The days of the week}

Solutions:

- a. {April, June, September, November}
- b. {2, 4, 6.....98}
- c. {Sunday, Monday, Tuesday, Wednesday, Thursday, Friday, Saturday}
- 3. Describe each set in words:
 - a. $A = \{a, b, c, d, e, f, g, h, i, j\}$
 - b. $B = \{2, 3, 5, 7, 11, 13, 17, \dots, 29\}$
 - c. C = {Divider, protractor, compass, ruler}

Solutions:

- a. A is the set of the first 10 letters of the alphabet.
- b. B is the set of prime numbers between 1 and 30.
- c. C is a set of mathematical instruments.
- 4. Let C be the set of countries in Africa. Insert either \in or \notin in the blank spaces:
 - a. Sierra Leone ____ C
 - b. Spain ____ C
 - c. United Kingdom _____ C

Solutions:

- a. Sierra Leone $\in C$
- b. Spain ∉ C
- c. United Kingdom $\notin C$
- 5. Complete the following:
 - a. If $D = \{$ the days of the week $\}$ find the cardinality of D.
 - b. If $Y = \{The months of the year that have 31 days\}$ find the cardinality of Y.
 - c. If $W = \{$ whole numbers greater than 1 and less than 10 $\}$ find n(W).

Solutions:

- a. D = {Sunday, Monday, Tuesday, Wednesday, Thursday, Friday, Saturday} There are 7 elements in the set, so n(D) = 7
- b. Y = { January, March, May, July, August, October, December} There are 7 elements in the set D, so n(Y) = 7
- c. $W = \{2, 3, 4, 5, 6, 7, 8, 9\}$ There are 8 elements in the set W, so n(W) = 8

Practice

1. Define each of the following sets of things by their common property:

- a. {London, Madrid, Moscow, Washington DC, Freetown}
- b. {Bee, Frog, Monkey, Wasp}
- c. $\{2, 4, 6, 8, 10, 12, 14, 16, 18\}$
- 2. List the elements of the following sets:
 - a. The set of months of the years with the first letter J
 - b. The set of odd numbers between 0 and 20
 - c. The set of even numbers less than 10
- 3. Describe the following sets in words:
 - a. {Freetown, Conakry, Monrovia, Abuja}
 - b. {Cow, chicken, pig, goat, sheep}
 - c. {Red, orange, yellow, green, blue, indigo, violet}
- 4. Make each of the statements true or false by replacing the symbol * by either \in

or ∉.

- a. 6 * {4, 9, 16, 25.....81}
- b. $\frac{4}{5}$ * {Whole numbers}
- c. $\sqrt{2} * \{\text{Irrational numbers}\}$
- 5. Complete the following:
 - a. If $D = \{ 4, 5, 6, 7, 8, 9 \}$ then n(D) =
 - b. If $X = \{1, 2, 3, 4, 5, 6\}$ then n(X) =
 - c. If $Y = \{0\}$ then n(Y) =

Lesson Title: Set-builder notation	Theme: Numbers and Numeration
Practice Activity: PHM1-L058	Class: SSS 1

By the end of the lesson, you will be able to write and interpret sets of values using set-builder notation.

Overview

Set-builder notation involves describing a set in terms of its characteristic or property, where only those with that specific property are considered elements of the set.

For example, consider $Q = \{x \mid x \text{ is a district in Sierra Leone}\}$. This statement says that "*Q* is the set of *x* such that *x* is a district in Sierra Leone". Here, the set Q is given by a property, instead of listing the members. The members of set Q are all of the districts in Sierra Leone. In this example, *x* is used as a symbol to represent a member of the set under consideration.

In set-builder notation, the vertical line | is read as "such that". You will sometimes see a colon (:) used instead of a vertical line.

Solved Examples

List the elements of the set: A = {x: x is multiples of 3 less than 20}
 Solution:

 $A = \{3, \, 6, \, 9, \, 12, \, 15, \, 18\}$

2. List the elements of the set $B = \{x | 1 \le x \le 5\}$ Solution:

 $B = \{1, 2, 3, 4, 5\}$

- Write in set-builder notation: D = {−4, −3, −2, −1....4}
 Solution: D = {x: x − 5 < x < 5}
- 4. Write in set-builder notation: E = {4, 8, 12, 16, 20, 24, 28} Solution:

 $E = \{x | x \text{ is multiples of 4 between 1 and 30} \}$

Practice

- 1. List the elements of the set: $A = \{x: x \text{ is factor of } 24\}$
- 2. List the elements of the set: $B = \{x: 10 \le x \le 15\}$

- 3. List the element of the set: $D = \{x: x \text{ is a prime number between 2 and 20}\}.$
- 4. List the elements of the set $B = \{x: x \text{ is a month of the year that has 31 days}\}$
- 5. Write in set-builder notation: $F = \{10, 100, 1,000, 10,000, ...\}$
- 6. Write in set-builder notation: $G = \{1, 3, 5, 7, 9\}$

Lesson Title: Finite and infinite sets	Theme: Numbers and Numeration
Practice Activity: PHM1-L059	Class: SSS 1

	Learning Outcome
	By the end of the lesson, you will be able to define and identify finite and infinite
set.	

Overview

A finite set is a set that comes to an end. For example, consider the set of even numbers less than 10. The elements of this set are {2, 4, 6, 8}. The members of a finite set are countable and can be listed.

An infinite set is uncountable, and the last element cannot be found. In an infinite set we add ellipses (...) at the end to show that the elements go on forever. For example, consider the set of all multiples of 3. There are an infinite number of multiples of 3, and they are {3, 6, 9, 12, 15, 18, 21...}.

Solved Examples

- 1. List the elements of each set, and state whether it is finite or infinite. Give the cardinality of each finite set.
 - a. $A = \{x | x \text{ is a factor of } 12\}$
 - b. $B = \{x: x \text{ is a multiple of 5}\}$
 - c. $C = \{x | x \text{ is a multiple of 6 less than 50} \}$
 - d. $D = \{x \mid x \text{ is a prime number}\}$
 - e. $E = \{x: x \text{ is an even number less than 30}\}$

Solutions:

- a. $A = \{1, 2, 3, 4, 6, 12\}$; finite; n(A) = 6
- b. $B = \{5, 10, 15, 20, 25, ...\}$; infinite
- c. $C = \{6, 12, 18, ..., 48\}$; finite; n(C) = 8
- d. $D = \{2, 3, 5, 7, 11, 13, ...\}$; infinite
- e. $E = \{2, 4, 6, 8, 10, ..., 28\}$; finite; n(E) = 14
- 2. State which of the sets is finite or infinite. List the elements of each.
 - a. Days of the week
 - b. The alphabet
 - c. Odd numbers
 - d. Whole numbers greater than 100
 - e. Prime numbers less than 35

Solutions:

- a. Finite; {Sunday, Monday, ..., Saturday}.
- b. Finite; {a, b, c, ..., z}
- c. Infinite; {1, 3, 5, 7, 9, …}

- d. Infinite; {101, 102, 103, 104, ...}
- e. Finite; {2, 3, 5, 7, ..., 31}

Practice

State whether each set is finite or infinite. List its members, using ellipses where necessary.

- 1. Factors of 18
- 2. Natural numbers
- 3. Months of the year
- 4. Multiples of 8 less than 70
- 5. Multiples of 7
- 6. Even numbers between 10 and 40
- 7. Prime numbers greater than 50

Lesson Title: Null/empty, unit and universal sets	Theme: Numbers and Numeration
Practice Activity: PHM1-L060	Class: SSS 1

By the end of the lesson, you will be able to define and identify null/empty sets, unit sets, and universal sets.

Overview

This lesson is on how to define and identify null/empty sets, unit sets and universal sets.

A set that does not contain any element is called a **null** or **empty set**. Such a set is represented by a curly bracket with nothing inside { } or the null symbol { \emptyset }. For example, the set A of cats with horns is a null set. We can write $A = \{\}$ or $A = \{\emptyset\}$.

A **unit set** is a set that has only one element, no more or less. Its cardinality is always one. For example, consider the set $F = \{x : x \text{ is a month with less than 30 days}\}$. We know that $F = \{February\}$, which is a unit set. The cardinality is n(F) = 1.

A **universal set** is the set of all elements or objects under consideration. For example, consider a school. We can use sets to discuss different groups of pupils in the school. For example, we have SSS1, SSS2, SSS3, and SSS4 pupils in the school. We have female and male pupils in the school. The universal set is the set of all pupils under consideration; in other words, all pupils in the school. A universal set is denoted by \mathcal{E} or U.

Solved Examples

- 1. Identify the sets below as empty/null sets or not:
 - a. $A = \{Goats with five legs\}$
 - b. $B = \{Months of the year with 45 days\}$
 - c. C = {Families with 2 children}

Solutions:

- a. A is an empty set.
- b. B is an empty set.
- c. C is not an empty set.
- 2. Decide whether the following sets are unit sets or not:
 - a. X = {Countries with capital city London}
 - b. $Y = \{x: x \text{ is the solution of } x + 1 = 2\}$
 - c. $Z = \{x: x \text{ is the value of } \sqrt{1}\}$
Solutions:

- a. X = {United Kingdom}.
 - n(X) = 1, so X is a unit set.
- b. Y = {1}; n(Y) = 1, so X is a unit set.
 c. Z = {+1, -1}
 - n(Z) = 2, so Z is not a unit set.
- 3. State the universal set of the following sets:
 - a. $C = \{Consonants\} and V = \{Vowels\}$
 - b. F = {Female Sierra Leoneans} and M= {Male Sierra Leoneans} **Solutions:**
 - a. $U = \{x: x \text{ is a letter of the alphabet}\}$

b. $U = \{ x : x \text{ is a Sierra Leonean} \}$

- 1. Decide whether each of the following sets are empty sets or not.
 - a. $G = \{Goats that can swim\}$
 - b. R = {Reptiles that can breast feed}
 - c. M = {Countries that have landed men on the moon}
 - d. P = {Female Presidents of Sierra Leone}
- 2. State the elements of each of the following unit sets:
 - a. D = {Month of the years with three letters}
 - b. $E = \{x: x \text{ is the solution of } x 2 = 5\}$
 - c. W = {The first day of the week}
- 3. State the universal set of the sets E = {*x*: *x* is an even number} and O = {*x*: is an odd number}
- 4. Given that $U = \{AII \text{ quadrilaterals}\}$, list the elements of the following:
 - a. C = {Cyclic quadrilaterals}
 - b. E = {Quadrilateral with two equal opposite sides}

Lesson Title: Equivalent and equal sets	Theme: Numbers and Numeration
Practice Activity: PHM1-L061	Class: SSS 1

Learning Outcome

By the end of the lesson, you will be able to define and identify equivalent and equal sets.

Overview

To be equivalent any pair of sets should have the same cardinality. In other words, sets are equivalent if they have the same number of elements. This implies that there is one- to-one correspondence between the elements of any two sets for them to be equivalent. For example, consider 2 sets, $A = \{1,2,3,4,5\}$ and $B = \{a, e, i, o, u\}$; n(A) = n(B) = 5. Thus, *A* and *B* are equivalent.

Note that all null/empty sets are equivalent. Similarly, all unit sets are equivalent.

Two sets are equal if and only if they have the same elements. This implies that for any pair of sets to be equal, they must first be equivalent and then have the same elements regardless of the order of occurrence of the elements. For example, if $P = \{1, 2, 3\}$ and $Q = \{2, 3, 1\}$ then P = Q. P and Q are equal sets.

Note that the order in which the members of a set are written does not matter in determining whether they are equal.

Solved Examples

- 1. Consider the following sets. Establish whether they are equivalent or not.
 - a. $P = \{ 5, 4, 3, 2, 1 \}$ and $Q = \{ J, U, L, Y, S \}$
 - b. $A = \{a, b, c, d, e\}$ and $R = \{m, n, o, p\}$
 - c. $F = \{\text{Hen, goat, cow, pig, cock}\}$ and $G = \{x: x > 5\}$

Solutions:

- a. Since n(P) = n(Q) = 5, P is equivalent to Q.
- b. n(A) = 5 and n(R) = 4. Since $n(A) \neq n(R)$ then set A is not equivalent to R.
- c. n(F) = 5 and n(G) is infinity so set F is not equivalent to G.
- 2. Consider the following sets and establish whether they are equal or not.
 - a. $W = \{x: 0 < x < 4\}$ and $V = \{x: 0 \le x \le 4\}$
 - b. C = {spoon, saucer, plate, pot}, and S = {pot, spoon, plate, saucer}
 - c. $D = \{x: x = \sqrt{4}\}$ and $E = \{x: x^2 4 = 0\}$
 - d. $X = \{ x: 2x = -16 \}$ and $Y = \{8\}$

Solutions:

a. $W = \{1, 2, 3\}$ and $V = \{0, 1, 2, 3, 4\}$, since the elements in W and V are not all identical, then $W \neq V$.

- b. The elements in C and S are identical, so C = S.
- c. Note that $D = \{+2, -2\}$ and $E = \{+2, -2\}$. D = E since both sets have identical elements
- d. Note that $X = \{-8\}$ and $Y = \{8\}$. Since $-8 \neq 8$, the elements in X and Y are not identical, so $X \neq Y$.

- 1. Consider the following pairs of sets. Establish whether they are equivalent or not.
 - a. $K = \{0, -1, 1, 5, 6\}$ and $L = \{\pi, \Phi, \rho, \gamma, \theta\}$
 - b. $A = \{u, v, w, x, y\}$ and $B = \{9, 10, 11\}$
 - c. C = {Toyota, Benz, Lexus, Nissan} and T = {Chrysler, Dodge, Pontiac}
 - d. N = {Lucy, Brima, Joe, Mary} and M = {x: 4 < x < 9}
- 2. Consider the following pairs of sets and establish whether they are equal or not.
 - a. $V = \{x : x > 0\}$ and $W = \{x : x < 0\}$
 - b. C = {January, February, March, April} and D = {December, November, October, September}
 - c. A = { x: x is an odd number between 0 and 10} and B = {1, 3, 5, 7, 9}
 - d. $X = \{x: x = \sqrt{25}\}$ and $Y = \{x: 5x 25 = 0\}$
 - e. $R = \{x: (x + 1)(x 1) = 0\}$ and $S = \{x: -1 \le x \le 1\}$
 - f. $I = \{1\}$ and $J = \{x : x 1 = 0\}$

Lesson Title: Subsets	Theme: Numbers and Numeration
Practice Activity: PHM1-L062	Class: SSS 1

	Learning Outcomes
	By the end of the lesson, you will be able to:
1.	Describe and identify the subsets of a given set.
2.	Represent subsets with Venn diagrams.
З	Lise correct symbols to demonstrate subsets

Overview

For any two sets A and B, if every element in set *B* is present in set *A*, then *B* is a subset of *A*.

A special notation is used to represent the relationships between subsets. In the case of A and B we say $B \subset A$. This is read as "set B is a subset of set A" or "B contained A". The symbol \subset means "is a subset of" or "is found in". The notation $\not\subset$ means "not a subset of".

If two sets are equal then, then they are also subsets of each other. For example, consider sets A = {1, 2, 3, 4} and B = {4, 3, 2, 1}. They have the same elements, so $B \subset A$ and $A \subset B$.

A Venn diagram is a diagram that shows the relationship between sets. It is most commonly drawn with circles and squares. A circle inside of another circle commonly shows subsets.

For example, the Venn diagram below shows $B \subset A$, where $A = \{1, 2, 3, 4, 5, 6, 7, 8\}$ and $B = \{3, 5, 6, 7\}$.



Note that the empty set is a subset of every set, and every set is a subset of itself.

Solved Examples

Given the set X = {m, n. o, p, q}, list any ten subsets of X.
 Solution:

Given X = {m, n, o, p, q}, ten subsets are: X₁ = {m, n, o, p}, X₂ = {m, n, o, q}, X₃ = {m, n, p, q}, X₄ = {m, o, p, q} $X_5 = \{n, o, p, q\}, X_6 = \{m, n\}, X_7 = \{m, o\}, X_8 = \{m, p\}, X_9 = \{m, q\}, X_{10} = \{m, n, o\}$

- 2. If $A = \{a, b, c, d, e, f, g, h\}$ identify whether the following sets are subsets of A:
 - a. $B = \{b, d, f, h\}$
 - b. $C = \{a, c, e, g\}$
 - c. $D = \{a, b, c, d, e\}$
 - d. E = {a, c, l, g}

Solutions:

- a. B is a subset of A; all elements of B are found in the set A.
- b. C is a subset of A; all the elements in set C are found in set A.
- c. D is a subset of A; all the elements in the set D are found in set A.
- d. E is not a subset of A; only the elements in E are found in A, I is not in A.
- 3. Given the set $M = \{1, 2, 3, 4, 5, 6, 7, 8\}$ which of the following sets are subsets of M?
 - a. i. N = {2, 4, 6, 8}
 - ii. P = {1, 2, 5, 7, 9}
 - iii. Q = {1, 2, 3, 4 5}

b. Draw Venn diagrams to illustrate the relationships between set M and each of its subsets in part a.

Solutions:

- a. i. N is a subset of M; all the elements in N are in M.
 - ii. P is not a subset of M; 9 is an element of p but not in M.
 - iii. Q is a subset of M; all the elements in Q are in M.
- b. Diagrams showing that N and Q are subsets of M:





List all the subsets of the set Z = {U, V, W}.
 Solution:

The subsets of $Z = \{U, V, W\}$ are: $Z_1 = \{U, V, W\}, Z_2 = \{U, V\}, Z_3 = \{U, W\}, Z_4 = \{V, W\}$ $Z_5 = \{U\}, Z_6 = \{V\}, Z_7 = \{W\}$ and $Z_8 = \{\}$.

Practice

- 1. Given the set $S = \{a, b, c, d\}$ list all its subsets.
- 2. Given the set $X = \{x: 0 \le x \le 20, where x \text{ is an integer}\}$, identify which of the following sets are subsets of X:
 - a. $W = \{x: 3 < x < 9 \text{ where } x \text{ is an integer}\}$
 - b. $V = \{x: 0 \le x \le 5, \text{ where } x \text{ is an integer}\}$
 - c. $T = \{x : x > 0\}$
- 3. Represent the relationship between the following set B and its subsets V and C on a Venn diagram.
 - $B = \{x: x \text{ is an alphabet between a and f} \}$

 $V = \{x: x \text{ is a vowel}\}$

- $C = \{x: x \text{ is a consonant}\}$
- 4. List all the subsets of the set $M = \{x: x \text{ is a multiple of } 2 \text{ up to } 6\}$

Lesson Title: Intersection of 2 sets	Theme: Numbers and numeration
Practice Activity: PHM1-T2-W16-L063	Class: SSS 1

))	Learning	0	utco	omes	,
11		-			

- By the end of the lesson, you will be able to:
- 1. Describe and identify the intersection of 2 sets.
- 2. Represent the intersection of two sets with Venn diagrams.
- 3. Use the correct symbols for intersection.

Overview

The intersection of two sets A and B is the set that contains all the elements of A that also belong to B (or all the elements of B that also belong to A) but no other elements. For example, take set A to be the colours of the flag of Sierra Leone: $A = \{green, white, blue\}$ and set B to be the colours of the flag of Ghana: $B = \{red, yellow, green\}$.

The intersection of set A and B is the element(s) that is/are common to both sets A and B. It is denoted by $A \cap B$ and is read as "A intersection B".

The intersection of two sets can be drawn using a Venn diagram. A Venn diagram is an important tool allowing relationships between sets to be visualised graphically. In the Venn diagrams below, the shaded portion shows the intersection of the two sets A and B.



In I we have $A \cap B$.

In II note that if $B \subset A$, then $B \cap A = B$.

Solved Examples

1. Consider the two sets A and B:

$$A = \{2, 4, 6, a, b\}$$

- a. Find $A \cap B$
- b. Draw a Venn diagram to illustrate both sets



Solutions:

a. A \cap B = {6, a}, because 6 and a are common to both sets A and B.

b. Venn diagram:



- 2. Consider two sets P and Q:
 - P = {multiples of 3 less than 13]
 - Q = {Odd numbers less than 10}
 - a. List the members of both sets P and Q.
 - b. Find $P \cap Q$.
 - c. Draw a Venn diagram to illustrate $P \cap Q$.

Solutions:

- a. $P = \{3, 6, 9, 12\}$ and $Q = \{1, 3, 5, 7, 9\}$
- b. $P \cap Q = \{3, 9\}$, because 3 and 9 are common to both sets.
- c. Venn diagram:



- 3. Set A is prime numbers less than 15, and set B is even numbers less than 15.
 - a. Find $A \cap B$.
 - b. Draw a Venn diagram to illustrate set A and set B.

Solutions:

- a. List the elements of set A and set B:
 - A = {prime numbers less than 15}
 - = {2, 3, 5, 7, 11, 13}
 - $B = \{even numbers less than 15\}$
 - $= \{2, 4, 6, 8, 10, 12, 14\}$

Therefore, we have $A \cap B = \{2\}$, because 2 is the only common element.

b. Venn diagram:



Practice

1. Consider two sets P and Q:

$$P = \{6, 12, 18, 24\}$$
$$Q = \{12, 24, 36, 48\}$$

- a. Find $P \cap Q$.
- b. Draw a Venn diagram to illustrate set P and set Q.
- 2. Consider two sets R and S:

R = {factors of 12}

- S = {prime numbers less than 4}
- a. List the elements of sets P and set S.
- b. Find $R \cap S$.
- c. Illustrate set R and set S in Venn diagram.
- 3. Consider two sets A and B:
 - A = {multiples of 4 less than 19}
 - B = {even numbers greater than 7 but less than 19}
 - a. List the elements of set A and set B.
 - b. Find $A \cap B$.
 - c. Illustrate set A and set B on a Venn diagram.
- 4. From the Venn diagram below:
 - a. List the elements of set A and set B.
 - b. Identify $A \cap B$.



Lesson Title: Intersection of 3 sets	Theme: Numbers and Numeration
Practice Activity: PHM1-L064	Class: SSS 1

	Learning Outcomes
	By the end of the lesson, you will be able to:
1.	Describe and identify the intersection of 3 sets.
2.	Represent the intersection of 3 sets with a Venn diagram.

Overview

This lesson is on identifying the intersection of 3 sets and representing them by a Venn diagram.

The intersection of three sets X, Y and Z is the set of elements that are common to sets X, Y and Z. It is denoted by $X \cap Y \cap Z$ which is read as "X intersection Y intersection Z".

The Venn diagram drawn below shows the intersection of 3 sets. The intersection $A \cap B \cap C$ is the shaded area in each diagram.



The elements of the sets can be shown in a Venn diagram. To draw a Venn diagram, start by filling in the elements in the intersection first. Work outwards, filling in the elements that are in only one set last.

Solved Examples

- 1. For the sets:
 - $A=\{10,\,11,\,12,\,13,\,14,\,15\}$
 - $B = \{10,\,12,\,13,\,15,\,16,\,17\}$
 - $C=\{11,\,12,\,13,\,14,\,15,\,17\}$
 - a. Find $A \cap B$, $B \cap C$, $A \cap C$, and $A \cap B \cap C$
 - b. Illustrate the sets on a Venn diagram

Solutions:

a. $A \cap B = \{10, 12, 13, 15\}$ $B \cap C = \{12, 13, 15, 17\}$ $A \cap C = \{11, 12, 13, 14, 15\}$ $A \cap B \cap C = \{12, 13, 15\}$ b.



- 2. For the sets:
 - $X = \{ 4, \, 6, \, a, \, b, \, 8 \} \\ Y = \{ 4, \, a, \, b, \, c, \, 5 \}$
 - $Z = \{5, a, b, d, 8, 9\}$
 - a. Find $X \cap Y$, $Y \cap Z$, $X \cap Z$, and $X \cap Y \cap Z$.
 - b. Illustrate your answer with a Venn diagram.

Solutions:

a.
$$X \cap Y = \{4, a, b\}$$

 $Y \cap Z = \{5, a, b\}$
 $X \cap Z = \{8, a, b\}$
 $X \cap Y \cap Z = \{a, b\}$

b.



- 3. U is the universal set consisting of all positive integers *x* such that {22 < *x* < 42}.
 P, Q and R are subsets of U such that P = {*x*: *x* prime numbers}, Q = {*x*: *x* is an odd numbers above 30} and R = {23, 31, 32, 37, 41}
 - a. Find $P \cap Q$, $Q \cap R$, $P \cap R$, $P \cap Q \cap R$.
 - b. Illustrate your answer in a Venn diagram.

Solutions:

First, list the elements of each set:

 $U = \{23,\,24,\,25,\,26,\,27,\,28,\,29,\,30,\,31,\,32,\,33,\,34,\,35,\,36,\,37,\,38,\,39,\,40,\,41\}$

 $\mathsf{P} = \{23,\,29,\,31,\,37,\,41\}$

Q={31,33,35,37,39,41}.

R = {23, 31, 32, 37, 39}

a. Find the intersections, $P \cap Q$, $Q \cap R$, $P \cap R$ and $P \cap Q \cap R$:

$$P \cap Q = \{31, 37, 41\}$$

 $Q \cap R = \{31, 37, 39\}$
 $P \cap R = \{23, 31, 37\}$
 $P \cap Q \cap R = \{31, 37\}$

b. Draw the Venn diagram:



- 1. Consider the following sets, where U is the universal set:
 - $U=\{2,\,4,\,6,\,8...20\}$
 - $A = \{multiples of 4 less than 20\}$
 - $B = \{even numbers less than 14\}$
 - $C = \{8, \, 12, \, 16, \, 18, \, 20\}$
 - a. Find: i. $A \cap B$; ii. $B \cap C$; iii. $A \cap C$, iv. $A \cap B \cap C$
 - b. Draw a Venn diagram to illustrate $A \cap B \cap C$
- 2. If U = {1, 2, 3, 4, 5, 6, 7} and P,Q, and R are subsets, such that P = {1, 3, 6},
 - $Q=\{3,\,4,\,6\}, and \ R=\{2,\,3,\,7\},$
 - a. Find, $P \cap Q$, $P \cap R$, $Q \cap R$, $P \cap Q \cap R$.
 - b. Illustrate the intersections with a Venn diagram.
- 3. M, L, and T are subsets of the universal set U, where $U = \{x: 1 \le x \le 15\}$ and x is an integer. M consists of odd numbers, L consists of even numbers and T consists of multiples of 3.
 - a. Find $M \cap L$, $M \cap T$, $L \cap T$, and $M \cap L \cap T$.
 - b. Draw a Venn diagram to illustrate U, M, L, and T.
- The Venn diagram shown is a universal set U with subsets X,Y, and Z. Use it to find: X∩Y, X∩Z, Y∩Z, X∩Y∩Y.



Lesson Title: Disjoint set	Theme: Numbers and Numeration
Practice Activity: PHM1-L065	Class: SSS 1

Learning	Outcomes
----------	----------

- By the end of the lesson, you will be able to:
- 1. Describe and identify disjoint sets.
- 2. Represent disjoint sets with Venn diagram.

Overview

This lesson is on describing and identifying disjoint sets. Any two sets are said to be disjoint if they have no elements in common. That is, their intersection is the empty/null set. In symbols, $A \cap B = \emptyset$ means that A and B are disjoint sets.

Disjoint sets A and B can be illustrated using a Venn diagram as shown below:



Events are often used in probability. In terms of events, disjoint events cannot happen at the same time (they are mutually exclusive).

The following are considered as disjoint events:

- (i) Laughing and whistling.
- (ii) A football game and a lawn tennis match on the same pitch.
- (iii) Going North and South at the same time.
- (iv) Day and night occurring at the same time.
- (v) Getting a head and a tail in a single toss of a fair coin.

Solved Examples

Given the sets, P = {a, e, i, o, u} and Q = {m, n, p, q, r} show that P and Q are disjoint sets and illustrate the two sets on a Venn diagram.
 Solution:

 $P = \{a, e, i, o, u\},\ Q = \{m, n, p, q, r\}$

 $P \cap Q = \Phi$



 Given the sets: M = {set of lawyers in Sierra Leone}, N = {Set of medical doctors in Sierra Leone}, establish that the sets M and N are disjoint sets and illustrate on a Venn diagram.

Solution:

$$\label{eq:main_star} \begin{split} &\mathsf{M} = \{ \texttt{set of lawyers in Sierra Leone} \} \\ &\mathsf{N} = \{ \texttt{Set of medical doctors in Sierra Leone} \} \\ &\mathsf{M} \cap \mathsf{N} = \Phi \end{split}$$



- 3. Given the set $X = \{x: -4 \le x < 0\}, Y = \{x: 4 \le x < 8\}$
 - a. List the elements of sets X and Y
 - b. Show that sets X and Y are disjoint sets
 - c. Show X and Y on a Venn diagram

Solutions:

- a. $X = \{x: -4 \le x < 0\} = \{-4, -3, -2, -1\}$ $Y = \{x: 4 \le x < 8\} = \{4, 5, 6, 7\}$
- b. Sets X and Y do not have any element in common $X \cap Y = \Phi$
- c. Venn diagram:



- 1. Given the sets M = {-1, -2, -3, -4} and N = { $\sqrt{2}$, π , $\sqrt{3}$, $\sqrt{5}$ } show that M and N are disjoint sets. Show them on a Venn diagram.
- Given W = {Breed of goat} and N = {Breed of chicken}, show that sets M and N are disjoint sets and show them on a Venn diagram.
- 3. Given sets $X = \{x: 3 \le x \le 6\}$ and $Y = \{x: -3 \le x \le 0\}$
 - a. List the elements of sets X and Y.
 - b. Establish whether sets X and Y are disjoint sets.
 - c. Show X and Y on a Venn diagram.
- 4. If set $P = \{i, j, k, l\}$ and $R = \{4, 6, 8, 10\}$ show that sets P and Q are disjoint sets and illustrate them on a Venn diagram.

Lesson Title: Union of two sets	Theme: Numbers and Numeration
Practice Activity: PHM1-L066	Class: SSS 1

	Learning Outcomes
	By the end of the lesson, you will be able to:
1.	Describe and identify the union of two sets.
2.	Represent the union of the two sets with a Venn diagram.
2	Lies the correct symbol for union

3. Use the correct symbol for union.

Overview

The union of any two sets A and B is the set that contains elements or objects that belong to either sets A or B or both.

The union of two sets A and B is the set formed by putting the two sets together.

The union of 2 sets A and B is written in symbols as $A \cup B$. It is read 'A union B'.

If a member appears in both sets, it is listed only once in the union. Also note that when the members of sets are numbers, they are listed in ascending order (smallest member first).

These 3 diagrams show the union of sets. The shaded portion shows the union of the two sets A and B.



Solved Examples

 Given the sets A = {orange, pineapple, apple, mango, banana} and B = {plate, spoon, knife, cup}; find the union of the two sets and show the union on a Venn diagram.

Solution:

 $A \cup B = \{$ plate, spoon, knife, cup, orange, pineapple, apple, mango, banana $\}$



- 2. Given the set $X = \{x: -2 < x < 2, x \text{ is an integer}\}$ and $Y = \{x: -2 \le x \le 2, x \text{ is an integer}\}$
 - a. List the elements of X and Y
 - b. Find the union of X and Y
 - c. Show the union of X and Y in a Venn diagram

Solutions:

- a. $X = \{x: -2 < x < 2; x \text{ is an integer}\} = \{-1, 0, 1\}$ $Y = \{x: -2 \le x \le 2, x \text{ is an integer}\} = \{-2, -1, 0, 1, 2\}$
- b. $X \cup Y = \{-2, -1, 0, 1, 2, \} = Y$, note that $X \subset Y$.
- c. Venn diagram:



- 3. Given the sets $V = \{a, e, i, o, u\}$ and $C = \{a, b, c, d, e\}$:
 - a. Find the union of V and C

b. Show the union of V and C in a Venn diagram **Solutions:**

- a. $V \cup C = \{a, b, c, d, e, i, o, u\}$
- b. Venn diagram:



4. Given the sets:

 $M = \{x: x \text{ is a prime number up to } 10\}$

- N = {x: x is an odd positive integer up to 10}
- $P = \{x: x \text{ is an even positive integer up to 10}\}$
- a. List the elements of sets M, N and P
- b. Find the union of sets M, N and P.
- c. Show the union of the sets in a Venn diagram

Solutions:

- a. List the elements of the sets:
 - M = {x: x is a prime number up to 10} = {2, 3, 5, 7}
 - N = {x: x is an odd positive integer up to 10} = {3, 5, 7, 9]
 - $P = \{x: x \text{ is an even positive integer up to } 10\} = \{2, 4, 6, 8, 10\}$
- b. $M \cup N \cup P = \{2, 3, 4, 5, 6, 7, 8, 9, 10\}$
- c. Venn diagram:



Practice

- 1. If M = {5, 7, 8, 10, 11} and N = {4, 5, 6, 7} write the set M \cup N. Show M and N on a Venn diagram.
- 2. Given $U = \{a, b, c, d, e, f\}$ and:
 - $\begin{aligned} X &= \{a, \, b, \, c, \, d\} \\ Y &= \{c, \, d, \, e\} \\ Z &= \{b, \, d, \, f\}, \end{aligned}$
 - a. List the members of the following: i. $X \cup Y$ ii. $Y \cup Z$ iii. $X \cup Z$
 - b. Show $X \cup Y$ and $X \cup Z$ in Venn diagrams.
- The members of a quiz team at a school for a competition form a universal set U where U = {Alice, Brima, David, Grace, Moses, Sama Yomba, Sia, Lamin, Kandeh}. If:
 - A = {members whose name end in **a**}
 - B = {Members with three letter names}
 - $C = \{members whose names begin with letter$ **S** $\}.$

List the following sets:

a. AUB b. BUC c. AUC

4. The figure below is a Venn diagram representing a Universal set U and the subsets X and Y.



List the elements of the following sets:

- a. X∪Y
- b. X∪U
- c. YuU

Lesson Title: Complement of a set	Theme: Numbers and Numeration
Practice Activity: PHM1-L067	Class: SSS 1

Learning Outcomes	
-------------------	--

- By the end of the lesson, you will be able to:
- 1. Describe and identify the complement of a set.
- 2. Represent the compliment of a set with a Venn diagram.

Overview

If U is the universal and A is a subset of U, then the complement of A is the set of members which belong to the universal set U but do not belong to A. In other words, the complement of a set is equal to elements in the universal set minus elements in the set.

The complement of a set A is written as A' and is read as "A prime". For example, if $U = \{3, 7, 9, 11, 13\}$, and $A = \{3, 9, 11\}$. Then A' = $\{7, 13\}$

The following are the properties of the complement of a set:

- The complement of the universal set is the empty set. Complement of U is the set which does not contain any element (U' = { }).
- The complement of an empty set is the universal set under consideration.
 (Ø' = U).
- The union of a set and its complement give the universal set $(A \cup A' = U)$.
- The intersection of a set and its complement is the empty set; there are no elements that exist in both sets (A ∩ A' = Ø).

The diagram below illustrate complements of some sets. They are shown in symbols, and how to read each set of symbols is given.



 $(A \cup B)'$ "A union B, all prime"



"B prime"



 $(A \cap B)'$ "A intersection B, all prime"



"prime of A subset B"

Solved Examples

- 1. If $U = \{-3 \le x < 3\}$ and $A = \{-1 \le x \le 2\}$, a. List the elements of U and A b. Find A' **Solutions:** a. $U = \{-3, -2, -1, 0, 1, 2\}$ $A = \{-1, 0, 1, 2\}$ b. A'= $\{-3, -2\}$
- 2. If A and B are subsets of the universal set U such that $A = \{1, 3, 4, 6\}$, $B = \{2, 4, 7, 8\}$, and $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$, list the elements of:

u

12

13

10

16

11

15

18

14

- a. A'
 b. B'
 c. A'∩B'
 Solutions:
 a. A' = {2, 5, 7, 8, 9, 10}
 b. B' = {1, 3, 5, 6, 9, 10}
- c. A'∩B' = {5, 9, 10}
- 3. Use the Venn diagram below to identify and list the following sets.
 - a. U, X, Y
 - b. X'
 - c. Y'
 - d. X'∩Y'

Solutions:

a. U = {11, 12, 13, 14, 15, 16, 17, 18, 19} X = {12, 13, 14, 18} Y = {11, 14, 16, 18} b. X' = {11, 15, 16, 17, 19} c. Y' = {12, 13, 15, 17, 19} d. X' \cap Y' = {15, 17, 19}

- 4. Use the Venn diagram below to find the following sets.
 - a. A'
 - b. B'
 - c. A'∩B'



Solutions:

First, identify the sets:

 $\label{eq:U} \begin{array}{l} U = \{k,\,l,\,m,\,n,\,p,\,q,\,r,\,s,\,t\} \\ A = \{k,\,p,\,q,\,r\} \\ B = \{l,\,m,\,q,\,r\} \\ a. \ A' = \{l,\,m,\,n,\,t\} \\ b. \ B' = \{k,\,p,\,n,\,t\} \\ c. \ A' \cap B' = \{n,\,t\} \end{array}$

- 1. U = {4 $\leq x \leq 12$ } and x is an integer. P is a subset of U, such that P = {6 $\leq x < 10$ }
 - a. List the element of sets U and P
 - b. Find P'
- 2. U = {10, 20, 30, 40, 50, 60}. A and B are subsets of U such that A = {20, 30, 60}, B = {20, 30, 50}, find:
 - a. A'
 - b. B'
 - c. A'∩B'
- 3. Using the Venn diagram below, identify and list the following sets:
 - a. U, P and Q
 - b. P'
 - c. Q'
 - d. P'∩Q'



- 4. Use the Venn diagram below to find the following sets:
 - a. U, M, and N
 - b. M'
 - c. N'
 - d. M'∩N'



Lesson Title: Problem solving with 2 sets	Theme: Numbers and Numeration
Practice Activity: PHM1-L068	Class: SSS 1

Learning Outcome

By the end of the lesson, you will be able to diagram and solve problems involving 2 sets, including real-life problems.

Overview

When we have two sets A and B then:

- $n(A \cup B)$ is the number of elements present in either of the sets A or B.
- $n(A \cap B)$ is the number of elements present in both the sets A and B.
- For two sets A and B: $n(A \cup B) = n(A) + n(B) n(A \cap B)$

This is one formula that is often used in problem solving with sets. The formula states that to find the total number of elements in A and B, then we add the number of elements in each set, and subtract the number of elements in their intersection. This is because we do not want to count the elements in their intersection twice.

Solved Examples

1. If A and B are two finite sets such that n(A) = 32, n(B) = 18 and $n(A \cup B) = 42$, find $n(A \cap B)$.

Solution:

$$\begin{split} n(A \cup B) &= n(A) + n(B) - n(A \cap B) \\ n(A \cap B) &= n(A) + n(B) - n(A \cup B) \\ &= 32 + 18 - 42 \\ &= 50 - 42 \\ &= 8 \end{split}$$

- 2. A test was conducted for a class of 70 pupils. Fifty passed Mathematics and 40 passed English. Each pupil passed at least one subject.
 - a. Illustrate this information on a Venn diagram.
 - b. How many pupils passed both Mathematics and English? **Solutions:**

Let $U = \{number of pupils that took the test\}$. Then, $n\{U\} = 70$

- $P = \{number of pupils who passed Mathematics\}, and n\{P\} = 50$
- Q ={number of pupils who passed English}, and n{Q} = 40
- x = number of pupils who passed both Maths and English = P \cap Q

 $n(x) = n(P \cap Q)$

a. The Venn diagram shows the above information. \rightarrow

Number of pupils who passed Mathematics only= 50 - x

Number of pupils who passed English only= 40 - xThis is shown in the Venn diagram. \rightarrow

b. To find $x = n(P \cap Q)$, add the number of elements in the three regions and equate it to n{U}. Solve the equation for *x*.

$$(50 - x) + x + (40 - x) = 70$$

$$50 - x + x + 40 - x = 70$$

$$50 + 40 - x = 70$$

$$90 - x = 70$$

$$x = 90 - 70$$

$$= 20$$

- 3. In a school with a total population of 320 pupils, 118 of them play football only and 102 play volleyball only. Each pupil plays at least one of the two games.
 - a. Draw a Venn diagram to illustrate this information.
 - b. How many pupils play:
 - i. Both games
 - ii. Football

Solutions:

Let $U = \{pupil \text{ population in the school}\}$. Then $n\{U\} = 320$.

- F = Set of pupils who play Football
- V =Set of pupils who play Volleyball
- n = number of pupils who play both games

Note: Here, do not subtract n from the other regions because of the word **only.**

- a. Venn diagram →
- b. i. Those who play both games = n. Add the 3 regions and equate to U, then solve for n.

$$\begin{array}{rcrr} 118+n+102 &=& 320\\ 220+n &=& 320 \end{array}$$

n = 320 - 220 = 100

ii. Number of pupils who play football

$$= n + 118$$

= 100 + 118
= 218







 In the Venn diagram, set P has 80 members, set Q has 60 members while there are members in P∩Q and 120 members in P∪Q. use the Venn diagram to answer the following questions



- a. Shade the region which shows P only
- b. Find the number of members in $P \cap Q$
- c. Find the number of members in Q only

Solutions:

a.



b. $P \cap Q = x$

b. Price = x Region for P only = 80 - xRegion for Q only = 60 - xAdd the three regions and equate to $n(P \cup Q) = 120$ 80 - x + 60 - x + x = 120 80 + 60 - x = 120 140 - x = 120 x = 140 - 120 = 20c. Members in Q only = 60 - x = 60 - 20 = 40

- 1. In a school of 120 pupils, 70 are in the Lawn tennis team and 80 are in the Volleyball team. Each pupil is in at least one team.
 - a. Illustrate this information on a Venn diagram.
 - b. Find how many pupils are in both teams.
 - c. Find how many pupils are in Lawn tennis team only.
- In a village of 180 adult inhabitants, 60 speaks only Mende fluently and 70 speaks only Kono fluently. Each adult speaks at least one of the two languages
 - a. Draw a Venn diagram to illustrate this information
 - b. How many adults speak: i. Both languages fluently; ii. Kono; iii. Mende
- 3. In a class of 49 pupils, 27 offer French and 32 offer Krio. Nine pupils do not offer any of the languages.
 - a. Draw a Venn diagram to illustrate this information.
 - b. How many stunts offer: i. Both languages; ii. Only French; iii. Only Krio

- 4. Use the Venn diagram below to answer the following questions:
 - a. Find the number of members in $P \cap Q$.
 - b. Find the number of members in:
 - i. P only
 - ii. Q only



Lesson Title: Problem solving with 3 sets – Part 1	Theme: Numbers and Numeration
Practice Activity: PHM1-L069	Class: SSS 1

Learning Outcome

By the end of the lesson, you will be able to diagram and solve problems involving 3 sets, including real-life problems.

Overview

If A, B and C are subsets of U the universal set then the Venn diagram below shows the various regions of a problem with 3 sets.

The following are elements in each section of the diagram.

- I is the intersection of all the sets. $(A \cap B \cap C)$
- Il is the elements of A and B only. $(C' \cap (B \cap A))$
- III is the elements of A and C only. $(B' \cap (A \cap C))$
- IV is the elements of B and C only. $(A' \cap (B \cap C))$
- V is the elements in A only. $(A \cap (B \cup C)' \text{ or } A \cap B' \cap C')$
- VI is the elements in B only. $(B \cap (C \cup A)' \text{ or } B \cap C' \cap A')$
- VII is the elements in C only. $(C \cap (B \cup A)' \text{ or } C \cap B' \cap A')$
- VIII is the elements that are not in any of the 3 sets.

This is the first of 2 lessons on solving problems involving 3 sets.

Solved Examples

- 1. Given three sets X, Y and Z such that $n(X \cap Y \cap Z) = 4$, $n(X \cap Y) = 9$, $n(X \cap Z) = 5$, $n(Y \cap Z) = 6$, $n(X \cap Y' \cap Z') = 7$, $n(X' \cap Y' \cap Z) = 2$ and $n(X' \cap Y \cap Z') = 4$,
 - a. Draw the Venn diagram
 - b. Find: i. n(X) ii. n(Y) iii. n(Z) **Solutions:**
 - a. See Venn diagram on the right.
 - b. i. Find n(X) by adding all elements in set X:

$$n(X) = 7 + (9 - 4) + 4 + (5 - 4)$$

= 7 + 5 + 4 + 1
= 17

ii. Find n(Y) by adding all elements in set Y:

$$n(Y) = 4 + (9 - 4) + (6 - 4) + 4$$

= 4 + 5 + 2 + 4
= 15
Find n(Z) by adding all elements in

iii. Find n(Z) by adding all elements in set Z:

$$n(Z) = 4 + (5 - 4) + 2 + (6 - 4)$$
$$= 4 + 1 + 2 + 2$$
$$= 9$$





- 2. There are 73 pupils in a class. Twenty of them study history, 18 study art and 16 study government. Ten study all three subjects, 15 study both history and government, 12 study both history and art, while 14 study both art and government.
 - a. Illustrate the information on a Venn diagram.
 - b. Using the Venn diagram determine the number of pupils who study at least two subjects.

Solutions:

Let the set of pupils in the class be U; so total number of pupils will be n(U). Let the set of pupils who study history be H this implies that n(H) = number of pupils who study history. Let the set of pupils who study art be A; so n(A) = number of pupils who study art. Let the set of pupils who study government be G; so n(G) = number of pupils who study government.

From the problem, we have n(U) = 73, n(H) = 20, n(A) = 16, n(G) = 16, $n(H \cap A \cap G) = 10$, $n(H \cap G) = n(H \cap A) = 12$ and $n(A \cap G) = 14$.

- a. See Venn diagram at right.
- b. The number of pupils that study at least two subjects will be represented by adding each segment in the intersections of the sets, as follows:



n(at least two subjects) = $n(H \cap A \cap G') + n(H \cap G \cap A') + n(A \cap G \cap H') + n(H \cap A \cap G)$ = (15 - 10) + (14 - 10) + (12 - 10) + 10= 5 + 4 + 2 + 10= 21

- A group of prospective voters were asked to choose at least one candidate of three, represented by the colours red, green and white. Twenty of them preferred green, 17 preferred red and 16 preferred white. Eight preferred a coalition of white and red, 9 preferred green and red 12 preferred green and white and 5 preferred all three colours.
 - a. Draw a Venn diagram to illustrate the data.
 - b. From your diagram find:
 - i. The total number of prospective voters surveyed.
 - ii. The total number of voters that preferred only one candidate (colour). **Solutions:**

All prospective voters prefer at least one candidate, therefore there is no complement. So:

Let U = {set of prospective voters}

- $G = \{$ those who prefer green $\}$, implying n(G) = 20
- $R = \{$ those who prefer red $\}$, implying n(R) = 17

W = {those who prefer white}, implying n(W) = 16

 $n(W \cap R) = 8$, $n(G \cap R) = 9$, $n(W \cap G) = 12$, $n(G \cap R \cap W) = 5$

- a. See Venn diagram at right.
- Let the number of those voters that prefer green only be g, those who prefer red only be r and those who prefer white only be w.

n(G) =
$$g + (9 - 5) + 5 + (12 - 5)$$

20 = $g + 4 + 5 + 7$
20 = $g + 16$
 $g = 20 - 16 = 4$
n(W) = $w + (8 - 5) + 5 + (12 - 5)$
16 = $w + 3 + 5 + 7$
16 = $w + 15$
 $w = 16 - 15 = 1$
n(R) = $r + (9 - 5) + 5 + (8 - 5)$
17 = $r + 4 + 5 + 3$
17 = $r + 12$
 $r = 17 - 12 = 5$
i. Total number of prospective voters:
n(GURUW) = $g + (9 - 5) + 5 + (12 - 5) + (8 - 5) + r + w$

= g + r + w = 4 + 5 + 1 = 10



Practice

 In a class of 45 pupils, it is known that 24 study Music, 20 study Chemistry and 22 study Biology. All the pupils study at least one subject of the three. Three do all three subjects while 7 do Music and Biology. Six do Music and Chemistry but not Biology and 8 do Chemistry and Biology. How many pupils study: a. Only one subject; b. Two subjects only.

= 4 + 4 + 5 + 7 + 3 + 5 + 1 = 19ii. Number of voters that preferred only one candidate:

- 2. At a Junior Secondary School there are 12 teachers. Of these, 6 teach Mathematics, 5 teach Integrated Science, and 4 teach Social Studies, 2 teach Mathematics and Integrated Science but no one teaches both Mathematics and Social Studies.
 - a. Draw a Venn diagram to illustrate the data.
 - b. Find: i. The number of teachers who teach Integrated Science and Social Studies; ii. The number of teachers who teach Social Studies only.

Lesson Title: Problem solving with 3 sets – Part 2	Theme: Numbers and Numeration
Practice Activity: PHM1-L070	Class: SSS 1

Learning Outcome

By the end of the lesson, you will be able to diagram and solve problems involving 3 sets, including real-life problems.

Overview

This is the second of 2 lessons on problem solving with 3 sets.

If 3 sets A, B and C are subsets of a universal set U, then the following rule holds:

 $n(U) = n(A \cup B \cup C) = n(A) + n(B) + n(C) - n(A \cap B) - n(A \cap C) - n(B \cap C) + n(A \cap B \cap C)$

This rule is used throughout this lesson.

Solved Examples

1. A group of people were asked whether they liked soccer, basketball or rugby. Their responses are as shown in the table below;

Game liked	No. of people
All three games	7
Soccer and rugby	11
Soccer and basketball	10
Basketball and rugby	9
Soccer	34
Rugby only	18
Basketball only	16
None of the three games	3

- a. Represent the data in a Venn diagram
- b. How many people liked soccer only?
- c. How many people were surveyed?
- d. How many people liked exactly two games?

Solutions:

Let U = set of people surveyed

- S = set of soccer lovers
- B = set of basketball lovers
- R = set of Rugby lovers

Then we have the following:

 $n(S \cap B \cap R) = 7 \qquad n(S) = 34$

- $n(S \cap R) = 11$ $n[R \cap (B \cup S)'] = 18$
- $n(S \cap B) = 10$ $n[B \cap (S \cup R)'] = 16$

 $n(B \cap R) = 9 \qquad n(B \cap R \cap S)' = 3$

- a. See Venn diagram on the right.
- b. Let the number of soccer lovers only be *x*, so that n[S∩(B∪R)'] = *x*.
 Create an equation using the total number of people who like soccer, and solve for *x*:
- n(S) = x + (10 7) + 7 + (11 7) 34 = x + 3 + 7 + 4 34 = x + 14x = 34 - 14 = 20



n(U) = x + (10 - 7) + 16 + (11 - 7) + 7 + (9 - 7) + 18 + 3n(U) = 20 + 3 + 16 + 4 + 7 + 2 + 18 + 3 = 73

d. To find the number of people who liked exactly two games, add segments that represent exactly 2 games:

$$= (10 - 7) + (11 - 7) + (9 - 7)$$
$$= 3 + 4 + 2 = 9$$

- 2. In a survey, 74 out of 88 tourists interviewed said they had visited at least the Freetown cotton tree (T), Lumley Beach (B) or old Fourah Bay College (F). Of these, 19 had visited only Lumley Beach and Cotton Tree, 30 visited Lumley Beach and old Fourah Bay College, 26 visited old Fourah Bay College and the cotton tree. No one had visited only the Cotton tree, 10 visited only Lumley Beach, 7 visited only Old Fourah Bay College and *x* had visited all three sites.
 - a. Draw a Venn diagram to illustrate the information.
 - b. Write a suitable equation in x and solve it.

c. Find i. $n[(B \cap T) \cup F]$ ii. $n[(B \cap F)' \cup T]$ iii. n(T)Solutions:

Note that: n(U) = 88, $n(T \cup B \cup F) = 74$, $n(B \cap T) = 19$, $n(B \cap F) = 30$, $n(F \cap T) = 26$, $n[T \cap (B \cup F)'] = 0$, $n[B \cap (T \cup F)'] = 10$, $n[F \cap (B \cup T)'] = 7$, and $n(B \cap T \cap F) = x$

a. Venn diagram:



b. $n(B\cup T\cup F) = 10 + (19 - x) + x + (30 - x) + 0 + (26 - x) + 7$ 74 = 10 + 19 - x + x + 30 - x + 26 - x + 7 (equation in x) 74 = 92 - 2x 2x = 92 - 74 $x = \frac{18}{2} = 9$



c. (i)
$$n[(B\cap T)\cup F] = [(19 - x) + x] + (30 - x + 26 - x + 7)$$

 $= [(19 - 9) + 9] + (30 - 9 + 26 - 9 + 7)$
 $= (10 + 9 + 21 + 17 + 7)$
 $= 64$
(ii) $n[(BUF)'\cap T] = 0$
(iii) $n(T) = 19 - x + 0 + x + 26 - x$
 $= 19 + 26 - x$
 $= 19 + 26 - 9$
 $= 36$

- In a sample survey of 47 people about the local languages they can speak in Sierra Leone, 31 were found to speak Krio, 26 Temne and 16 Mende. Three were found to speak all three languages, 2 spoke Temne and Mende only, and 6 spoke Mende only.
 - a. Illustrate the information on a Venn diagram.
 - b. Find the number of people who speak:
 - i. Temne only; ii. Only one language; iii. Only two languages
- There are 70 women in a club. Each plays at least one of the following games: Ludo (L), cards (C) and dice (D). If 20 play Ludo and an equal number play two games only:
 - a. Illustrate the information on a Venn diagram.
 - b. Find the number of women who play Ludo.
- 3. The Leone star coach invited 20 players for a soccer match. Eight of them were attackers, 12 of them were defenders and 12 were midfielders. The coach realised that 5 could play in the attack and in midfield, 6 defence and midfield, 2 only in attack and 3 could play all three roles.
 - a. Illustrate the information on a Venn diagram.
 - b. Find the number of players that can play:
 - i. Only midfield; ii. Exactly two roles

Lesson Title: Use of variables	Theme: Algebraic Processes
Practice Activity: PHM1-L071	Class: SSS 1

2	Lea	rning	Οι	Jt	cor	nes
///	_	-		-		

- By the end of the lesson, you will be able to:
- 1. Identify that variables represent unknown numbers.
- 2. Identify the values of variables in simple algebraic equations (e.g. 2 + x = 5).

Overview

This lesson is on how to identify the values of variables in simple algebraic expressions.

A variable is a letter that is used to represent a number. For example, consider the expression 2 + x = 5. The variable *x* represents a specific number that will sum to 5 when 2 is added to it.

While x is the most common variable used in algebra, any letter can be used.

An algebraic expression is a collection of letters and symbols combined with at least one or more of the mathematical operations, +, -, × or ÷. For example, these are some algebraic expressions: 3m + n - 5, $5pq^3$, and $3pst \div 9p^2t^2s$

The **coefficient** of an algebraic term is the number in front of a variable. For example, in 8m, 8 is the coefficient of m. The coefficient is multiplied by the variable, so that $8m = 8 \times m$.

A number on its own (without any variable) is called a constant term in an algebraic expression. For example, in 3x + 5, 5 is a constant term.

Solved Examples

1. Identify the variables in the algebraic expressions:

a. $5\pi r^2 h$ b. x + y - 5 c. v = u + 9.8t

Solutions:

- a. r and h are variables. Recall that π has a known value, which is 3.14....
- b. x and y are variables
- c. u, v and t are variables
- 2. Identify the coefficients of x and y and the constant term in the following algebraic expressions:
 - a. -6x + 5y + 7 b. $\frac{6}{7}y x 1$ c. x + y + 10

Solutions:

- a. -6x + 5y + 7: The coefficient of x is -6The coefficient of y is 5 The constant term is 7
- b. $\frac{6y}{7} x 1$: The coefficient of *x* is -1 The coefficient of *y* is $\frac{6}{7}$ The constant term is -1 c. x + y + 10:
- The coefficient of x is 1
 The coefficient of y is 1
 The constant term is 10
- 3. Write the following statements as algebraic expressions:
 - a. 8 less than a certain number
 - b. 7 more than a certain number
 - c. Half a certain number
 - d. y times a certain number
 - e. 3 less than 2 times a certain number

Solutions:

Let *x* be the "certain number" in each example. Then we have:

- a. *x* 8
- b. *x* + 7
- c. $\frac{x}{2} \text{ or } \frac{1}{2}x \text{ or } 0.5x$
- d. $y \times x = xy$
- e. 2x 3
- 4. Find the value of the variable in each of the following expressions:
 - a. x 2 = 3
 - b. 2y = 10
 - c. 3a = 6
 - d. 3 + b = 7

Solutions:

- a. x = 5, because 5 2 = 3
- b. y = 5, because $2 \times 5 = 10$
- c. a = 2, because $3 \times 2 = 6$
- d. b = 4, because 3 + 4 = 7

- 1. Identify the variables in the algebraic expressions:
 - a. $\frac{4}{3}\pi r^3$
 - b. m + n 4

c.
$$x = ut + \frac{1}{2}at^2$$

- 2. Identify the coefficients and constants in each expression:
 - a. -5a + 6y 10
 - b. $\frac{4}{5}y + 7 x$
 - c. x y + 10
- 3. Write the following statements as algebraic expressions:
 - a. A certain number times 60.
 - b. 100 times as big as y
 - c. y more than two times x
- 4. Find the value of the variable in each of the following expressions:
 - a. x + 1 = 14
 - b. 4a = 16
 - c. 2b = 7 1
 - d. z 4 = 2

Lesson Title: Simplification – grouping terms	Theme: Algebraic Processes
Practice Activity: PHM1-L072	Class: SSS 1

Learning Outcome

By the end of the lesson, you will be able to simplify algebraic expressions by grouping like terms.

Overview

In algebra, like terms are terms that have the same variable, and the variables have the same power. For example, 2a, 3a, 5a are all like terms, with the variable a to the power 1. As another example, $5p^2$ and $8p^2$ are like terms.

Like terms can be combined by adding or subtracting to give a single term. The result will have the same variable with the same power. The coefficients of the terms are added or subtracted. For example, consider these combinations:

a + a = (1 + 1)a = 2aa + a + a = (1 + 1 + 1)a = 3a4a - a = (4 - 1)a = 3a

Unlike terms are two or more terms that are not like terms. That is, they do not have the same variables or powers. Examples of unlike terms are:

$$2a + 3b + 4c$$

 $3m^2 + 2m^3$

Unlike terms cannot be combined.

Solved Examples

1. Simplify: 8x - 2y - 4x + 5ySolution:

8x - 2y - 4x + 5y	=	8x - 4x + 5y - 2y	Collect like terms
	=	(8-4)x + (5-2)y	Combine like terms
	=	4x + 3y	

2. Simplify: 7p - 3q + 3p + 2qSolution:

$$7p - 3q + 3p + 2q = 7p + 3p + 2q - 3q$$
 Collect like terms
= $(7 + 3)p + (2 - 3)q$ Combine like terms
= $10p - q$

3. Simplify the following algebraic expressions:

```
a. 12e + 5f - 4e - 2f
      b. 2m + 5n - 3m - 4n
      c. 11x - 10y - 10x + 12y
      d. 3u - 3 + 4v - 2u + 7 - 2v
      Solutions:
      a.
       12e + 5f - 4e - 2f = 12e - 4e + 5f - 2f Collect like terms
                          = (12-4)e + (5-2)f Combine like terms
                          = 8e + 3f
      b.
       2m + 5n - 3m - 4n = 2m - 3m + 5n - 4n Collect like terms
                          = (2-3)m + (5-4)n Combine like terms
                          = -m + n
      C.
      11x - 10y - 10x + 12y = 11x - 10x + 12y - 10y
                             = (11-10)x + (12-10)y
                             = x + 2y
      d.
       3u - 3 + 4v - 2u + 7 - 2v = 3u - 2u + 4v - 2v + 7 - 3
                                 = (3-2)u + (4-2)v + (7-3)
                                 = u + 2v + 4
4. Simplify the following algebraic expressions:
      a. pq + 12xp + 5pq - 3xp
      b. a + 3ab + 6 + 7a - 2ab - 1
      Solutions:
      a.
      pq + 12xp + 5pq - 3xp = pq + 5pq + 12xp - 3xp
                             = (1+5)pq + (12-3)xp
                             = 6pq + 9xp
      b.
       a + 3ab + 6 + 7a - 2ab - 1
                                = a + 7a + 3ab - 2ab + 6 - 1
                                  = (1+7)a + (3-2)ab + (6-1)
```

65

= 8a + ab + 5

Practice

Simplify the following algebraic expressions:

- 1. 4y 3x + 5x 3y2. 9a + 4b - 11a + 3b3. 6m + 11n - 4m + 2n - m + n4. 2xy - 4xp + 3xy + 3xp5. 8mn + 9m + 4mn - 10m6. 4pq - 5pq + 8qr + 4qr + 3pq - 4qr7. $12p^2q - 4pq^2 + pq^2 - 4p^2q$ 8. 3xyz - 4xz + 6xyz + 3xz
- 9. 4ab + 7 3a 3 2ab + 7a
| Lesson Title: Simplification – removing brackets | Theme: Algebraic processes |
|---|----------------------------|
| Practice Activity: PHM1-L073 | Class: SSS 1 |

By the end of the lesson, you will be able to simplify algebraic expressions by removing brackets.

Overview

In today's lesson, we are looking at algebraic expressions that contain brackets. For example, 2(5x - 4). In order to simplify such expressions, we must first remove the brackets. In removing brackets, multiply the term outside the bracket by each of the terms inside the bracket.

We must be very careful with signs when removing brackets. When there is a (+) sign before the bracket, the sign inside the brackets does not change when the brackets are removed. When there is a negative sign (-) in front of the brackets, the signs inside the bracket change when the brackets are removed. This is because of the rules of multiplication. Remember that multiplying a negative by a positive gives a negative. Multiplying a negative and a negative gives a positive.

After removing brackets, collect any like terms together and combine them.

Solved Examples

1.	Simplify: $2(5x - $ Solution:	- 4)	
	2(5x - 4)	= (2)(5x) + (2)(-4) = 10x - 8	Multiply each term by 2
2.	Simplify: $-4(2y)$ Solution: -4(2y-3)	= (-4)(2y) + (-4)(-3) = -8y + 12	3) Multiply each term by –4

3. Simplify: 8p - 3(2q + 2p) **Solution:** 8p - 3(2q + 2p) = 8p + (-3)(2q) + (-3)(2p) Remove the brackets = 8p - 6q - 6p = 8p - 6p - 6q Collect like terms = 2p - 6q 4. Simplify: 5n - 3(2m + 3n)Solution: 5n - 3(2m + 3n) = -3(2m + 3n)

5n - 3(2m + 3n) = 5n - 6m - 9n Remove the brackets = 5n - 9n - 6m Collect like terms = -4n - 6m

5. Remove the brackets and simplify 3(y + 5) + 6(y + 2). **Solution:**

3(y+5) + 6(y+2) = 3y + 15 + 6y + 12 Remove the brackets = 3y + 6y + 15 + 12 Collect like terms = 9y + 27

6. Remove the brackets and simplify 7(2p + 3) - 4(p - 2). **Solution:**

7(2p+3) - 4(p-2) = 14p + 21 - 4p + 8 Remove the brackets = 14p - 4p + 21 + 8 Collect like terms = 10p + 29

7. Remove the brackets and simplify 4(b-2a) + 3(a-3b) - 4(b-a). Solution:

$$4(b-2a) + 3(a-3b) - 4(b-a) = 4b - 8a + 3a - 9b - 4b + 4a$$

= -8a + 3a + 4a + 4b - 9b - 4b
= -a - 9b

Practice

Remove brackets and simplify the following algebraic expressions:

- 1. 5(x-4)
- 2. -7(3y 4)
- 3. 9m 2(m + n)
- 4. 11u 3u(2v + 3)
- 5. -(2x+4y)-(x-y)
- 6. 8(-3m+2n) 2(m+n)

Lesson Title: Simplification – expanding brackets	Theme: Algebraic Processes
Practice Activity: PHM1-L074	Class: SSS 1

By the end of the lesson, you will be able to simplify algebraic expressions by expanding brackets.

Overview

In expanding an expression containing brackets, we rewrite the expression in an equivalent form without any brackets. Expanding brackets is the same as removing brackets. For example, to expand 3(x + 2) means we multiply each of the terms inside the bracket by 3. Thus, 3(x + 2) = 3x + 6.

This lesson is on expanding brackets when there are 2 sets of brackets, for example, (x + 5)(x + 10). In this expression, 2 binomials are multiplied. Binomials are algebraic expressions with 2 terms. Multiply each term in one bracket by each term in the other bracket. After removing the brackets, always collect like terms together and simplify.

Solved Examples

1.	Expand $(x + 5)(x + 1)$ Solution:	.0)	
	(x+5)(x+10)	= x(x + 10) + 5(x + 10)	Multiply each term of the first binomial by the second binomial
		$= x^2 + 10x + 5x + 50$	Remove the brackets
		$= x^2 + 15x + 50$	Combine like terms
2.	Expand $(y-4)(y-3)$	3)	
	Solution:		
	(y-4)(y-3)	= y(y-3) - 4(y-3)	Multiply
		$= y^2 - 3y - 4y + 12$	Remove the brackets
		$= y^2 - 7y + 12$	Combine like terms
3.	Expand $(3a + b)(2a + b)$	- <i>b</i>)	
	Solution:		
	(3a + b)(2a - b)	= 3a(2a-b) + b(2a-b)	 Multiply
		$= 6a^2 - 3ab + 2ab - b^2$	Remove the brackets
		$= 6a^2 - ab - b^2$	Combine like terms

4. Expand $(x - 5)^2$

Solution:

$(x-5)^2$	= (x-5)(x-5)	Rewrite
	= x(x-5) - 5(x-5)	Multiply
	$= x^2 - 5x - 5x + 25$	Remove the brackets
	$= x^2 - 10x + 25$	Combine like terms

5. Expand
$$(3mn - 4)(2mn + 3)$$

Solution:
 $(3mn - 4)(2mn + 3) = 3mn(2mn + 3) - 4(2mn + 3)$ Multiply
 $= 6m^2n^2 + 9mn - 8mn - 12$ Remove the brackets
 $= 6m^2n^2 + mn - 12$ Combine like terms

6. Expand and simplify (5m + 6n)(u - v) + (3m + 2n)(2u + v)Solution:

Step 1. Expand the first two brackets:

(5m + 6n)(u - v) = u(5m + 6n) - v(5m + 6n) Multiply = 5mu + 6nu - 5mv - 6nv Remove the brackets **Step 2.** Expand the last two brackets:

$$(3m+2n)(2u+v) = 2u(3m+2n) + v(3m+2n)$$
 Multiply
= $6mu + 4nu + 3mv + 2nv$ Remove the brackets

Step 3. Collect like terms after the two expansions:

5mu + 6mu + 6nu + 4nu - 5mv + 3mv - 6nv + 2nv= 11mu + 10nu - 2mv - 4nv

7. Expand and simplify $2p(2p-4)^2$

Solution:

Step 1. Expand the brackets:

$$(2p-4)^2 = (2p-4)(2p-4)$$

= 2p(2p-4) - 4(2p-4)
= 4p^2 - 8p - 8p + 16
= 4p^2 - 16p + 16

Step 2. Multiply by 2p:

 $\begin{array}{rl} 2p(2p-4)^2 &= 2p(4p^2-16p+16) \\ &= 8p^3-32p^2+32p \end{array}$

Practice

Expand and simplify the following:

1. (7x + 2)(x + 4)2. (p - 5)(p - 6)3. (m + 3)(m - 2)4. $(x - 4)^2$ 5. $(n + 3)^2$ 6. (3m + 2)(n + 3) + (5m - 4)(n + 2)7. $3x(x - 2)^2$ 8. $4(x + 6)^2$

Lesson Title: Factoring - common factors	Theme: Algebraic Processes
Practice Activity: PHM1-L075	Class: SSS 1

By the end of the lesson, you will be able to factorise algebraic expressions by determining common factors.

Overview

The focus of this lesson is how to factorise algebraic expressions by determining common factors. Factorising in algebra is the reverse of expanding.

Consider 2(a + b) = 2a + 2b. You have expanded 2(a + b) by removing brackets.

If you switch sides in the above equation you will get 2a + 2b = 2(a + b). We can factorise 2a + 2b by taking the common factor 2 out of each term in 2a + 2b to get 2 multiplied by (a + b).

The general method used to factorise terms with common factors in algebraic expressions is to find the greatest common factor (GCF) to all terms and bring it outside the bracket.

Variables can also be common factors. If every term of an expression has the same variable, it can be factorised. Remember that factoring is the same as dividing. If you factor a variable from an expression, subtract the exponents.

If you choose a factor that is not the GCF, you might factor a number or variable, and then realise there is more that you can factor. It is fine to factorise an expression more than once, just continue with the next factor.

To check for the correctness of your factorisation, you can always expand the factorised expression. If you get the original expression, then you are right. Otherwise, check your factorisation again.

Solved Examples

1. Factorise the expression 10a - 15b + 5Solution:

Notice that 5 is a common factor in each term of the expression. That is, each can be divided by 5: $\frac{10a}{5} = 2a$, $\frac{15b}{5} = 3b$, $\frac{5}{5} = 1$ Factor 5 from each term:

10a - 15b + 5 = 5(2a) - 5(3b) + 5(1)= 5(2a - 3b + 1)

2. Factorise: 4a - ab

Solution:

a is a common factor of each term. Take *a* out of the bracket:

4a - ab = a(4 - b)

3. Factorise $:8x^3 + 4x^2 + 2x$

Solution:

2x is a common factor of each term: $8x^{3} + 4x^{2} + 2x = 2x(4x^{2} + 2x + 1)$

4. Factorise $2a^7 + 3a^6 + a^5 + 4a^3$ Solution:

> a^3 is a common factor. $2a^7 + 3a^6 + a^5 + 4a^3 = a^3(2a^4 + 3a^3 + a^2 + 4)$

- 5. Factorise:
 - a. $64x^3 + 16x^2$
 - b. $5a^4 10a^3 + 15ab$

Solutions:

- a. $16x^2$ is a common factor: $64x^3 + 16x^2 = 16x^2(4x + 1)$
- b. 5a is a common factor: $5a^4 10a^3 + 15ab = 5a(a^3 2a^2 + 3b)$
- 6. Factorise:
 - a. $15p^{6}q^{4} 6p^{4}q^{3}$ b. $3xy^{3} - 3xy^{2} + 6x^{2}y^{2}$ **Solutions:** a. $3p^{4}q^{3}$ is a common factor: $15p^{6}q^{4} - 6p^{4}q^{3} = 3p^{4}q^{3}(5p^{2}q - 2)$ b. $3xy^{2}$ is a common factor:
 - $3xy^3 3xy^2 + 6x^2y^2 = 3xy^2(y + 2x 1)$

Practice

Factorise the following algebraic expressions:

- 1. 9*ab* 27
- 2. $x^2 4x$
- 3. $a^3 + ab$
- 4. $16x^4 8x^3 + 4x^2$
- 5. $81a^3 + 9a^2$
- 6. $12x^4 36x^3 + 48x^2$
- 7. $18s^5t^6 6s^4t^3 + 3s^2t^2$
- 8. $5uv^3 25uv^2 + 15u^2v^2$

Lesson Title: Factoring – grouping	Theme: Algebraic processes
Practice Activity: PHM1-L076	Class: SSS 1

By the end of the lesson, you will be able to factorise algebraic expressions by grouping common terms.

Overview

This lesson focuses on factorising algebraic expression by grouping terms with common factors. We often have expressions with four terms which can be factored by grouping the terms that have common factors.

For example, consider the algebraic expression: $x^2 - ax + bx - ab$

Notice that the 4 terms do not have 1 common factor. The absence of a common factor to all four terms does not mean the expression cannot be factorised. There are common factors to some pairs of terms which can be grouped for factorisation.

Note that you can change the order of the terms based on which terms have common factors. For the example expression, we can form a group with a common factor of x, and another group with a common factor of b. We then factor x and b.

Form the two groups: $x^2 - ax + bx - ab = (x^2 - ax) + (bx - ab)$

Factor x and b: x(x - a) + b(x - a)

We now have 2 terms in the expression, x(x - a) and b(x - a). Notice that (x - a) is a common factor in these terms. It can be factored out of the expression, leaving (x + b):

$$x^{2} - ax + bx - ab = x(x - a) + b(x - a) = (x - a)(x + b)$$

Solved Examples

1. Factorise completely: $2a^3 + 4a^2 - 3a - 6$ Solution:

$$2a^{3} + 4a^{2} - 3a - 6 = (2a^{3} + 4a^{2}) + (-3a - 6)$$
 Create smaller groups
= $2a^{2}(a + 2) - 3(a + 2)$ Factor out each GCF
= $(a + 2)(2a^{2} - 3)$ Factor out $(a + 2)$

2. Factorise completely: $2x^3 - 6x^2 + 10x - 30$ Solution:

In this case, all 4 terms have a common factor, 2. When there are common factors, we should first factor them out. We can then proceed with the steps described above on the 4 terms inside brackets.

$$2x^{3} - 6x^{2} + 10x - 30 = 2(x^{3} - 3x^{2} + 5x - 15)$$

$$= 2[(x^{3} - 3x^{2}) + (5x - 15)]$$

$$= 2[x^{2}(x - 3) + 5(x - 3)]$$

$$= 2(x - 3)(x^{2} + 5)$$

Factor out each GCF
Factor out (x - 3)

3. Factorise completely: $18mn + 4m^2 - 27pn - 6pm$ **Solution:**

$$\begin{array}{ll} 18mn+4m^2-27pn-6pm &= 18mn-27pn+4m^2-6pm & \mbox{Reposition terms} \\ &= (18mn-27pn)+(4m^2-6pm) & \mbox{Create groups} \\ &= 9n(2m-3p)+2m(2m-3p) & \mbox{Factor GCFs} \\ &= (2m-3p)(9n+2m) & \mbox{Factor } (2m-3p) \end{array}$$

4. Factorise completely: xz + wy + xy + wzSolution:

xz + wy + xy + wz	= xz + wz + wy + xy	Reposition terms
	= (xz + wz) + (wy + xy)	Create groups
	= z(x+w) + y(x+w)	Factor out GCFs
	=(x+w)(z+y)	Factor out $(x + w)$

5. Factorise the expression $4uz - 6uv - 12vz + 2u^2$ Solution:

$$4uz - 6uv - 12vz + 2u^{2} = 4uq + 2u^{2} - 6uv - 12vz$$

$$= (4uq + 2u^{2}) + (-6uv - 12vz)$$

$$= 2u(2z + u) - 6v(2z + u)$$

$$= (2z + u)(2u - 6v)$$
Reposition terms
Create groups
Factor out GCFs
Factor out (2z + u)

Practice

Factorise the following expressions completely:

1.
$$ax - a + x - 1$$

2. $12eg - 4eh - 6fg + 2fh$
3. $cd - ce - d^2 + de$
4. $p + q + 5ap + 5aq$
5. $x^2 + 2x - 5x - 10$
6. $6a^2 - 3a + 4a - 2$
7. $x^2 + 2x + 3x + 6$
8. $3x - 2dy + 3y - 2dx$

Lesson Title: Substitution of values	Theme: Algebraic Processes
Practice Activity: PHM1-L077	Class: SSS 1

	Learning Outcome
	By the end of the lesson, you will be able to substitute values into given
algeb	praic expressions.

Overview

This lesson is on substituting values into given algebraic expressions.

In substitution, variables are replaced by numbers, and we evaluate the expression to find a final result. Substitution is a useful tool in algebra to find a value or to rewrite equations in terms of a single variable.

In substitution, you will need to observe the following rules:

- When substituting negative numbers put them in brackets () so that you can get the calculation right.
- Two like signs (-)(-) or (+)(+) become a positive sign and two unlike signs (+)(-) or (-)(+) become a negative sign. This follows the rule for multiplying negative numbers.
- Remember to use the correct order of operations (BODMAS).

Solved Examples

- 1. Find the value of x 2 if x = 6. Solution: We can substitute 6 for x. Instead of x - 2, we will have 6 - 2. Answer: 6 - 2 = 4.
- 2. If x = 3 and y = 4, what is x + xy? Solution:

Remember that two variables written together in a term (as in xy) means they are multiplied together.

 $\begin{array}{ll} x + xy &= 3 + (3)(4) & \qquad \text{Substitute 3 for } x \text{ and 4 for } y \\ &= 3 + 12 & \qquad \text{Simplify} \\ &= 15 & \qquad \end{array}$

3. Evaluate the expression $x^2 + xy$ given x = 3 and y = 4Solution:

$$x^{2} + xy = 3^{2} + (3)(4)$$

Substitute 3 for x and 4 for y
$$= 9 + 12$$

Simplify
$$= 21$$

4. Evaluate $x^2 + 2x + 1$ given x = 5. Solution:

 $x^{2} + 2x + 1 = 5^{2} + 2(5) + 1$ Substitute 5 for x = 25 + 10 + 1 Simplify = 36

5. Evaluate
$$2 - a + a^2$$
 given $a = -2$
Solution:

 $2-a+a^2 = 2-(-2)+(-2)^2$ Substitute -2 for a = 2+2+4 Simplify = 8

6. Evaluate $\frac{2y-z}{2x^2yz}$ when x = 1, y = -4 and z = -10Solution:

$$\frac{2y-z}{2x^2yz} = \frac{2(-4)-(-10)}{2(1)^2(-4)(-10)}$$

Substitute $x = 1, y = -4, z = -10$
$$= \frac{-8+10}{2(1)(40)}$$

Simplify
$$= \frac{2}{2\times40}$$

$$= \frac{2}{80} = \frac{1}{40}$$

7. Find the value of x in the equation $x = ut + \frac{1}{2}gt^2$, given u = 2, t = 5 and g = 10. Solution:

$$x = ut + \frac{1}{2}gt^{2}$$

Substitute $u = 2, t = 5 \text{ and } g = 10$
$$= 2(5) + \frac{1}{2}(10)(5)^{2}$$

Simplify
$$= 10 + 5(25)$$

$$= 10 + 125$$

$$= 135$$

Practice

- 1. What is the value of $x^2y y^2x$ if x = 3 and y = -1?
- 2. If a = -7 and b = 3, calculate the value of:

a.
$$\left(\frac{a+b}{a-b}\right)^2$$

b. $2a^2b + b^2a$
c. $27a^2 \div 12b^2$

- 3. What is the value of $\frac{str}{st-tr}$ if s = -3, t = 2 and r = -22?
- 4. Evaluate the value of C from the equation $C = \frac{5}{9}(F 32)$ given:
 - a. F = 59
 b. F = 95

Lesson Title: Addition of algebraic fractions	Theme: Algebraic processes
Practice Activity: PHM1-L078	Class: SSS 1

By the end of the lesson, you will be able to add algebraic expressions.

Overview

Recall the rules for adding fractions. Fractions with the same denominator can be added by simply adding the numerators. When fractions have different denominators, we determine the LCM of the denominators and change the denominators to their LCM before adding.

This lesson is on the addition of algebraic fractions. Algebraic fractions are fractions with a variable term in the numerator, in the denominator, or in both. Examples of algebraic fractions are: $\frac{x}{4}$, $\frac{2}{x+1}$, $\frac{x}{2+x}$. This lesson is on the addition of simple algebraic fractions, with variables in the denominators.

To add algebraic fractions, you will follow similar rules to adding numerical fractions. To add algebraic fractions, follow these steps:

- Find the lowest common multiple (LCM) of the denominators.
- Express all fractions with the LCM in the denominator.
- Once the fractions have the same denominator, they are **like fractions** and can be added.
- Add the fractions, and combine like terms if possible.
- If the answer can be simplified, simplify it.

Solved Examples

1. Simplify: $\frac{x}{3} + \frac{x}{2}$

Solution:

Find the LCM of the denominators (3 and 2). Their LCM is 6, so we will change the denominator of each fraction to 6 before adding.

$$\frac{x}{3} + \frac{x}{2} = \frac{2 \times x}{6} + \frac{3 \times x}{6}$$

$$= \frac{2x}{6} + \frac{3x}{6}$$

$$= \frac{2x + 3x}{6}$$

$$= \frac{2x + 3x}{6}$$

$$= \frac{5x}{6}$$

Express denominators as the LCM
Simplify the numerators
Add the numerators

2. Simplify: $\frac{2x}{9} + \frac{5x}{6}$

Solution:

Find the LCM of the denominators, which is 18.

 $\frac{2x}{9} + \frac{5x}{6} = \frac{2(2x)}{18} + \frac{3(5x)}{18}$ $= \frac{4x}{18} + \frac{15x}{18}$ $= \frac{4x + 15x}{18}$ $= \frac{4x + 15x}{18}$ $= \frac{19x}{18}$ Express denominators as the LCM Simplify the numerators Add

3. Simplify: $\frac{4x-1}{5} + \frac{2x+2}{5}$

Solution:

The denominators are the same, so proceed with addition:

$$\frac{4x-1}{5} + \frac{2x+2}{5} = \frac{4x-1+2x+2}{5}$$
 Add the numerators
$$= \frac{4x+2x+2-1}{5}$$
 Collect like terms
$$= \frac{6x+1}{5}$$
 Combine like terms

4. Simplify:
$$\frac{4x+3}{12} + \frac{2x+5}{8}$$

Solution:

Find the LCM of their denominators, which is 24.

$$\frac{4x+3}{12} + \frac{2x+5}{8} = \frac{2(4x+3)}{24} + \frac{3(2x+5)}{24}$$
Express the denominators as the LCM

$$= \frac{8x+6+6x+15}{24}$$
Simplify and add the numerators

$$= \frac{8x+6x+6+15}{24}$$
Collect like terms

$$= \frac{14x+21}{24}$$
Combine like terms

like terms

5. Simplify: $\frac{3x+6}{24} + \frac{2x-1}{8}$

Solution:

Find the LCM of their denominators, which is 24.

$\frac{3x+6}{24} + \frac{2x-1}{8}$	$=\frac{3x+6}{24}+\frac{3(2x-1)}{24}$	Express the denominators as the LCM
24 0	$=\frac{3x+6+3(2x-1)}{3x+6+3(2x-1)}$	Add
	$=\frac{3x+6+6x-3}{24}$	Simplify
	$=\frac{3x+6x+6-3}{24}$	Collect like terms
	$=\frac{9x+3}{24}$	Combine like terms
	$=\frac{3(3x+1)}{3(3x+1)}$	Factor the numerator
	$=\frac{\frac{24}{3x+1}}{8}$	Simplify

Practice

1. Express as a single fraction:

a.
$$\frac{a}{5} + \frac{4a}{5}$$

b. $\frac{2m}{9} + \frac{5m}{9}$
c. $\frac{5x}{7} + \frac{3x}{7}$
d. $\frac{x}{2} + 1$

2. Add:

a.
$$\frac{a}{5} + \frac{a}{3}$$

b. $\frac{x}{4} + x$
c. $\frac{n}{10} + \frac{n}{1}$
d. $\frac{x}{4} + \frac{x}{8}$

3. Simplify:

a.
$$\frac{3a+1}{4} + \frac{a-1}{3}$$

b. $\frac{2n+2}{7} + \frac{3n}{4}$
c. $\frac{3x}{9} + \frac{x-1}{6}$
d. $\frac{x+1}{5} + \frac{x}{10}$

Lesson Title: Subtraction of algebraic fractions	Theme: Algebraic fractions
Practice Activity: PHM1-L079	Class: SSS 1

By the end of the lesson, you will be able to subtract algebraic fractions.

Overview

This lesson is on subtraction of algebraic fractions. Recall that in the addition of algebraic fractions we find the LCM of the denominator, express the fraction in terms of the LCM and simplify to get a single fraction as the answer. We will follow similar steps to subtract algebraic fractions.

To subtract algebraic fractions, follow the steps:

- Find the lowest common multiple (LCM) of the denominators.
- Express all fractions with the LCM in the denominator.
- Once the fractions have the same denominator, they are **like fractions** and can be added or subtracted.
- Subtract the fractions. Subtract each of the like terms in the numerator.
- If the answer can be simplified, simplify it.

Be careful of a minus sign before a bracket. Where a minus sign occurs before a bracket, the signs of the terms inside the bracket change. For example, -(a - b) becomes -a + b. The negative sign is distributed to each term inside the brackets.

Solved Examples

1. Simplify: $\frac{3x}{4} - \frac{x}{3}$

Solution:

Find the LCM of the denominators, which is 12.

$\frac{3x}{4} - \frac{x}{3}$	$\frac{x}{3} =$	$\frac{3(3x)}{12} - \frac{4x}{12}$	Express denominators as the LCM
	=	$\frac{9x}{12} - \frac{4x}{12}$	Simplify numerator
	=	$\frac{9x-4x}{12}$	Subtract the numerators
	=	$\frac{5x}{12}$	Combine like terms

2. Simplify: $\frac{4a+1}{5} - \frac{4}{5}$

Solution:

The denominators are the same, so proceed with subtraction:

$$\frac{\frac{4a+1}{5} - \frac{4}{5}}{=} \frac{\frac{4a+1-4}{5}}{\frac{4a-3}{5}}$$

3. Simplify: $\frac{a-4}{2} - \frac{a-2}{6}$

Solution:

Find the LCM of the denominators, which is 6.

$\frac{a-4}{2} - \frac{a-2}{6}$	=	$\frac{3(a-4)}{(a-4)} - \frac{a-2}{(a-4)}$	Express denominators as the LCM
2 6	=	$\frac{3a-12}{6} - \frac{a-2}{6}$	Simplify the numerator
	=	3a-12-(a-2)	Subtract the numerators
	=	$\frac{3a-12-a+2}{6}$	Distribute the negative sign
	=	$\frac{3a-a-12+2}{\epsilon}$	Collect like terms
	=	$\frac{2a-10}{6}$	Combine like terms
	=	$\frac{2(a-5)}{2(3)}$	Factor out 2
	=	$\frac{a-5}{3}$	Simplify

4. Simplify: $\frac{m+4}{3} - \frac{2-m}{5}$

Solution:

Find the LCM of the denominators, which is 15.

 $\frac{m+4}{3} - \frac{2-m}{5} = \frac{5(m+4)}{15} - \frac{3(2-m)}{15}$ Express denominators as the LCM $= \frac{5(m+4) - 3(2-m)}{15}$ $= \frac{5m+20-6+3m}{15}$ $= \frac{5m+3m+20-6}{15}$ Collect like terms in the numerator and simplify $= \frac{8m+14}{15}$ $= \frac{2(4m+7)}{15}$

5. Simplify: $\frac{8x-8}{6} - \frac{2x-5}{7}$

Solution:

First, notice that we can simplify the fraction on the left. We can factor 2 from the numerator and denominator.

$$\frac{8x-8}{6} - \frac{2x-5}{7} = \frac{2(4x-4)}{2(3)} - \frac{2x-5}{7}$$
Factor the left fraction
$$= \frac{4x-4}{3} - \frac{2x-5}{7}$$
Simplify
$$= \frac{7(4x-4)}{21} - \frac{3(2x-5)}{21}$$
Express denominators as the LCM
$$= \frac{7(4x-4)-3(2x-5)}{21}$$
Subtract the numerators
$$= \frac{28x-28-6x+15}{21}$$
Simplify
$$= \frac{28x-6x-28+15}{21}$$
Collect like terms

$$=\frac{22x-13}{21}$$

Practice

1. Simplify:

a.
$$\frac{3x}{4} - \frac{2x}{4}$$

b. $\frac{2y}{3} - 1$
c. $\frac{2x+8}{5} - \frac{7}{5}$
d. $\frac{a}{3} - \frac{4a}{3}$
btract:

2. Subtract

a.
$$\frac{a}{3} - \frac{a}{4}$$

b. $\frac{4x}{5} - x$
c. $\frac{5x}{2} - \frac{2x}{3}$
d. $\frac{x}{5} - \frac{x}{4}$

Express as a single fraction in its lowest terms:

a.
$$\frac{5y}{2} - \frac{2y+6}{6}$$

b. $\frac{a-4}{2} - \frac{a-3}{3}$
c. $\frac{2y+2}{3} - \frac{y-1}{4}$
d. $\frac{2y+3}{3} - \frac{7}{12}$

Lesson Title: Linear equations	Theme: Algebraic processes
Practice Activity: PHM1-L080	Class: SSS 1

By the end of the lesson, you will be able to solve linear equations using the balance method.

Overview

A linear equation in one variable is an equation that can be written in the form ax + b = c where a, b and c are real numbers, x is the variable and $a \neq 0$. The highest power of the variable in a linear equation is 1, so linear equations are also called first-degree equations.

To use the balancing method, we solve by applying the same operation to both sides of an equal to sign. Our goal is to get the variable by itself.

These are possible steps in using the balancing method:

- Add the same quantity to each side.
- Subtract the same quantity from each side.
- Multiply each side by the same quantity.
- Divide each side by the same quantity.

You often need to perform multiple operations to get the answer. When balancing, perform addition or subtraction before multiplication and division.

Solved Examples

1. Solve for x in the equation x + 5 = 7. Solution: x + 5 = 7x + 5 - 5 = 7 - 5Subtract 5 from both sides x + 0 = 2x = 22. Solve 8 = 3x - 7. Solution: 8 = 3x - 78 + 7 = 3x - 7 + 7Add 7 to both sides 15 = 3x $= \frac{3x}{3}$ 15 Divide both sides by 3 3 3 5 = x

3. Solve 5y = 40.

Solution:

$$5y = 40$$

$$\frac{5y}{5} = \frac{40}{5}$$

$$y = 8$$

Divide both sides by 5

4. Solve for x if 5x - 3 = 3x + 7. Solution: 5x - 3 = 3x + 7 5x - 3 - 3x = 3x + 7 - 3x Su 2x - 3 = 72x - 3 + 3 = 7 + 3 Ac

 $\begin{array}{rcl} 2x & = & 10\\ \frac{2x}{2} & = & \frac{10}{2} \end{array}$

x = 5

Subtract 3x from both sides Add 3 to both sides

Divide both sides by 2

5. Solve 7y = 2y + 20. **Solution:**

$$7y = 2y + 20$$

$$7y - 2y = 2y - 2y + 20$$
 Sub

$$5y = 20$$

$$\frac{5y}{5} = \frac{20}{5}$$
 Division

$$y = 4$$

Subtract 2y from both sides Divide both sides by 5

6. Solve
$$4 - 7y = 3y + 4$$
. **Solution:**

$$4 - 7y = 3y + 4$$

$$4 - 7y - 3y = 3y - 3y + 4$$
Subtract 3y from both sides
$$4 - 10y = 4$$
Subtract 4 from both sides
$$-10y = 0$$

$$\frac{-10y}{-10} = \frac{0}{-10}$$
Divide both sides by-10
$$y = 0$$

Practice

Solve the following equations:

1. 11 + 5m = -42. 9y + 1 = 7y3. 3n - 15 = 45 4. 5y - 2 = 8y - 75. x + 4 = 7 - 2x6. 0 = 10 - 8y7. 3 - 4x = 5x + 128. 1 + 7m = 5m + 19. 5y + 6 = 3y - 5

Lesson Title: Linear equations with brackets	Theme: Algebraic Processes
Practice Activity: PHM1-L081	Class: SSS 1

By the end of the lesson, you will be able to solve linear equations that contain brackets.

Overview

This lesson is on solving linear equations that contain brackets.

To solve the equation containing brackets, first remove the brackets. Then balance the equation, as in the previous lesson.

Remember that when there is a negative sign outside the bracket, the signs inside the brackets change.

Solved Examples

1.	Solve $2(p + 1) = 18$. Solution:			
	2(p+1) =	18		
	2p + 2 =	18	Removi	ing the brackets
	2p + 2 - 2 =	18 – 2	Subtrac	ct 2 from both sides
	2p =	16		
	2p	16	Divide b	both sides by 2
	$\frac{1}{2}$ =	2		·
	p =	8		
2.	Solve $3(m + 6) = 2(m + 6)$	n - 3).		
	Solution:			
	3(m+6) =	2(m-3)		
	3m + 18 =	2m - 6		Remove brackets
	3m - 2m + 18 =	2m — 2m	- 6	Subtract $2m$ from both sides
	m + 18 =	-6		
	m + 18 - 18 =	-6 - 18		Subtract 18 from both sides
	m =	-24		
3.	Solve $5v - 3 = 3(v + 1)$	- 4).		
•	Solution:	-)-		
	5v-3	= 3(v + 4)	4)	
	5y - 3	= 3v + 1	2	Remove the brackets
	5v - 3v - 3	= 3v - 3	v + 12	Subtract 3v from both sides
	0, 0, 0	0, 0,	, , , ,	

$$2y = 12$$

$$\frac{2y}{2} = \frac{12}{2}$$

$$y = 6$$
Divide both sides by 2
$$y = 6$$

$$4. \text{ Solve } 8m - 4(m + 3) = 2m + 6.$$
Solution:
$$8m = 2m + 6$$

$$-4(m + 3)$$

$$8m - 4m - 12 = 2m + 6$$

$$4m - 12 = 2m + 6$$

$$4m - 12 = 2m - 6$$

$$2m - 12 = 12 = 2m - 6$$
Subtract $2m$ from both sides
$$2m - 12 = 6$$

$$2m - 12 + 12 = 6 + 12$$
Add 12 to both sides by 2
$$2m = 18$$

$$\frac{2m}{2} = \frac{18}{2}$$
Divide both sides by 2
$$m = 9$$

$$5. \text{ Solve } 5(y - 4) - 4(2y + 1) = 0$$
Solution:
$$5(y - 4) - 4(2y + 1) = 0$$
Solution:
$$5(y - 4) - 4(2y + 1) = 0$$
Collect like terms
$$-3y - 24 = 0$$

$$-3y - 24 + 24 = 0 + 24$$

$$\frac{-3y}{-3} = \frac{24}{-3}$$
Divide both sides by -3
$$y = -8$$

$$6. \text{ Solve } 2(3m + 5) - 3(m - 4) = 1$$
Solution:
$$2(3m + 5) - 3(m - 4) = 1$$

$$3m + 12 - 12$$
Subtract 12 from both sides
$$3m = -11$$

$$\frac{3m}{3} = -\frac{11}{3}$$
Divide both sides by 3
$$3m = -3\frac{2}{3}$$

Practice

Solve the following equations:

1. 2(a + 1) = 122. 2(m + 4) = 93. 4(x + 7) + 12 = 04. 3(y + 2) = 2 - y5. 4(2a - 7) = 3(4a - 9)6. 3(6 + 7x) + 2(1 - 5x) = 427. 4(3 - 5p) - 7(5 - 4p) + 3 = 0

Lesson Title: Linear equations with fractions	Theme: Algebraic Processes
Practice Activity: PHM1-L082	Class: SSS 1

By the end of the lesson, you will be able to solve linear equations that contain fractions.

Overview

This lesson is on solving linear equations that contain fractions, for example: $\frac{x}{6} = 8$.

When an equation contains a fraction, we can multiply to cancel the denominator. Remember to multiply both sides of the equation by the same number.

When there is more than 1 fraction in an equation, they can be cleared by multiplying throughout by the LCM of the denominators. After multiplying by the LCM, follow the steps from the previous 2 lessons. Remove any brackets, and balance the equation to solve for the variable.

Remember that you can check your solution by substituting the value of the variable into the equation. If both sides of the equation are equal, then the solution is correct.

Solved Examples

1. Solve
$$\frac{x}{6} = 8$$

Solution:

$$\frac{x}{6} = 8$$

$$6\left(\frac{x}{6}\right) = 6(8)$$
 Multiply by 6

$$x = 48$$

2. Solve: $\frac{x}{2} + 3 = 17$ Solution:

$$\frac{x}{2} + 3 = 17$$

$$2\left(\frac{x}{2}\right) + 2(3) = 2(17)$$
 Multiply by 2

$$x + 6 = 34$$

$$x + 6 - 6 = 34 - 6$$
 Subtract 6

$$x = 28$$

3. Solve for
$$x: \frac{x}{6} + \frac{x}{4} = 10$$

Solution:

$$\frac{x}{6} + \frac{x}{4} = 10$$

$$12\left(\frac{x}{6}\right) + 12\left(\frac{x}{4}\right) = 12(10)$$
Multiply each term by the LCM, 12
$$2x + 3x = 120$$
Simplify
$$5x = 120$$
Combine like terms
$$\frac{5x}{5} = \frac{120}{5}$$
Divide both sides by 5
$$x = 24$$

4. Solve the equation for $x: \frac{7}{24} = \frac{x}{8} + \frac{1}{6}$ Solution:

$$\frac{7}{24} = \frac{x}{8} + \frac{1}{6}$$

$$24\left(\frac{7}{24}\right) = 24\left(\frac{x}{8}\right) + 24\left(\frac{1}{6}\right)$$

$$7 = 3x + 4$$

$$7 - 4 = 3x + 4 - 4$$

$$3 = 3x$$

$$\frac{3}{3} = \frac{3x}{3}$$

$$1 = x$$

Multiply by 24, the LCM

Subtract 4

5. Solve for y in the equation:
$$\frac{3}{4}y - 2\frac{1}{3} = \frac{2}{3}y$$

Solution:

$$\frac{3}{4}y - 2\frac{1}{3} = \frac{2}{3}y$$

$$\frac{3y}{4} - \frac{7}{3} = \frac{2y}{3}$$
Conver
$$12\left(\frac{3y}{4}\right) - 12\left(\frac{7}{3}\right) = 12\left(\frac{2y}{3}\right)$$
Multiply
$$9y - 28 + 28 = 8y + 28$$
Add 28
$$9y = 8y + 28$$

$$9y - 8y = 8y - 8y + 28$$
Subtract
$$y = 28$$

rt mixed to improper fraction y each term by the LCM, 12

ct 8y

6. Solve for x:
$$\frac{x-1}{3} - \frac{x+1}{10} = 5$$

Solution:
 $\frac{x-1}{3} - \frac{x+1}{10} = 5$
 $30\left(\frac{x-1}{3}\right) - 30\left(\frac{x+1}{10}\right) = 30(5)$ Multiply each term by the LCM, 30
 $10(x-1) - 3(x+1) = 150$
 $10x - 10 - 3x - 3 = 150$ Remove the brackets (expand)
 $7x - 13 = 150$ Combine like terms
 $7x - 13 + 13 = 150 + 13$ Add 13

$$7x = 163$$

 $\frac{7x}{7} = \frac{163}{7}$ Divide by 7
 $x = 23\frac{2}{7}$

Practice

Solve the following equations:

1.
$$\frac{1}{5}x = 3$$

2. $\frac{2a}{5} + 1 = 9$
3. $\frac{y}{3} + \frac{y}{2} = 4$
4. $\frac{1}{5}x + \frac{1}{4}x = \frac{1}{2}$
5. $\frac{2}{3}(x - 6) = 8$
6. $\frac{k-3}{3} - \frac{k+2}{5} = 3$
7. $\frac{2p+7}{6} = \frac{2p-1}{3}$

Lesson Title: Word problems	Theme: Algebraic Processes
Practice Activity: PHM1-L083	Class: SSS 1

By the end of the lesson, you will be able to create and solve equations from word problems.

Overview

This lesson is on solving equations derived from word (story) problems. There are many types of word problems which involve relations among known and unknown numbers. These can be written in the form of equations.

To solve word problems you will need the following steps:

- Read the problem carefully and note what is given and what is required.
- Assign a variable to represent each unknown.
- Identify any value that is multiplied by the variable (the coefficient).
- Identify any value that is constant.
- Write the algebraic expression representing the situation.
- Solve the equation for the unknown variable.
- Check whether the answer satisfies the conditions of the problem.

Solved Examples

1. The sum of two numbers is 25. One of the numbers exceeds the other by 9. Find the numbers.

Solution:

Let one of the numbers be y. Then the other number is y + 9.

The sum of the two numbers is 25. Therefore, in this situation, y + y + 9 = 25. Solve the equation for *y*:

$$y + y + 9 = 25$$

$$2y + 9 = 25$$

$$2y + 9 - 9 = 25 - 9$$
 Subtract 9 from both sides

$$2y = 16$$

$$\frac{2y}{2} = \frac{16}{2}$$
 Divide both sides by 2

$$y = 8$$

We can find the other number using the formula: y + 9 = 8 + 9 = 17

Therefore, the two numbers are 8 and 17.

2. The difference between two numbers is 10. If one of the numbers is 4, what is the other number?

Solution:

Let the other number be y. The difference between 4 and y can either be 4 - y or y - 4.

If the difference is 10, then either:

$$\begin{array}{ll} 4 - y = 10 & y - 4 = 10 \\ 4 - 10 = y & \mathsf{OR} & y = 10 + 4 \\ -6 = y & y = 14 \\ y = -6 \end{array}$$

3. A girl is 5 years old and her mother is 27 years old. In how many years' time will the daughter be half her mother's age?

Solution:

Age of girl now = 5

Age of mother now = 27

Let x years be the number of years in which daughter will be $\frac{1}{2}$ her mother's age.

Age of daughter in x years = (5 + x)

Age of mother in x years = (27 + x)

For the daughter's age to be $\frac{1}{2}$ her mother's age, we will have $(5 + x) = \frac{1}{2}(27 + x)$. Solve the equation for x:

 $\frac{1}{2}(27 + x)$. Solve the equation for x:

 $(5+x) = \frac{1}{2}(27+x)$ 2(5+x) = 27+x 10+2x = 27+x 2x-x = 27-10 x = 17 yearsMultiply throughout by 2 Collect like terms

In 17 years' time, the girl will be half her mother's age.

4. One farmer has 119 chickens and another has 73. After they each sell the same number of chickens, the first farmer is left with three times as many chickens as the other. How many chickens did each sell? **Solution:**

If each farmer sells y chickens, the first farmer will be left write (119 - y) chickens and the second farmer will be left with (73 - y) chickens.

If the first farmer is left with 3 times that of the second after selling *y* chickens, then we have the equation (119 - y) = 3(73 - y). Solve for *y* to find how many chickens they each sold:

(119 - y) = 3(73 - y) 119 - y = 219 - 3y 3y - y = 219 - 119 2y = 100 $y = \frac{100}{2}$ y = 50 chickens

 $\begin{array}{rcl} 119-y &= 219-3y \\ 3y-y &= 219-119 \end{array}$ Distribute 3 on the right-hand side Collect like terms

- 5. The sum of 6 and one-third of a certain number is one less than twice the number.
 - a. Express this statement in algebraic terms.
 - b. What is the value of the number?

Solution:

a. Let the number be x. The sum of 6 and one-third of x is $(6 + \frac{x}{3})$. Twice the number x is 2x. One less twice x is (2x - 1). Therefore, we have the equation $6 + \frac{x}{3} = 2x - 1$.

b. Solve for *x*:

$$6 + \frac{x}{3} = 2x - 1$$

$$3\left(6 + \frac{x}{3}\right) = 3(2x - 1)$$

$$18 + x = 6x - 3$$

$$18 + 3 = 6x - x$$

$$21 = 5x$$

$$x = \frac{21}{5} = 4\frac{1}{5}$$

Multiply throughout by 3 to eliminate the fraction
Collect like terms

Practice

- 1. The result of adding 15 to a certain number and dividing the answer by 4 is the same as taking the number from 80.
 - a. Express this statement as an algebraic equation.
 - b. Find the value of the number.
- 2. A given number of matches are needed to fill 20 match boxes with the same number of matches in each box.
 - a. How many matches are there in each box?
 - b. If each box has three less match sticks, there will be enough sticks for 32 boxes. What is the total number of match sticks?
- 3. Bockarie is 11 years older than Kallon. In 5 years, Bockarie will be twice as old as Kallon. Find their present ages.
- 4. Sia and Kumba share Le7350.00 between them such that Sia gets Le950.00 more than Kumba. Find out how much money each gets.
- 5. Divide 59 mangoes into two parts so that one part is 7 mangoes less than five times the other part. How many mangoes are in the two parts?

Lesson Title: Substitution in formulae	Theme: Algebraic Processes
Practice Activity: PHM1-L084	Class: SSS 1

By the end of the lesson, you will be able to substitute given values into	a
formula.	

Overview

This lesson is on substituting given values into a formula. A formula is an equation in which variables represent quantities. For example, we have a formula for finding the perimeter P, of a rectangle. For a rectangle with length *l* and width *w*, perimeter is given by the formula P = 2(l + w), where *l* and *w* are in the same units. In this example, P is the subject of the formula. By substituting values of *l* and *w* into the formula, the corresponding value of P can be found.

Solved Examples

Ρ

1. Calculate the perimeter of a rectangle with a length of 8 cm and width of 6 cm. **Solution:**

= 2(l + w)	Formula
= 2(8 cm + 6 cm)	Substitute $l = 8$ and $w = 6$
= 2(14 cm)	Simplify
= 28 cm	

1. Given that a = 3, b = -2, find the value of the algebraic expression 2a + b. Solution:

Substitute the given values into the formula and simplify:

$$2a + b = 2(3) + (-2) = 6 - 2 = 4$$

2. Given that m = -5, n = 4, find the value of the algebraic expression $\frac{2m-n}{m^2}$. Solution:

Substitute the given values into the formula and simplify:

$$\frac{2m-n}{m^2} = \frac{2(-5)-(4)}{(-5)^2} = \frac{-10-4}{25} = \frac{-14}{25}$$

3. The circumference *C* of a circle is given by the formula $C = 2\pi r$, where *r* is the radius and π (pi) is a constant. Calculate the circumference of a circle with a radius of 28 cm. Use $\frac{22}{7}$ for π .

Solution:

С	$=2\pi r$	Formula
	$= 2 \times \frac{22}{7} \times 28$ cm	Substitute $\pi = \frac{22}{7}$ and $r = 28$

$$= 2 \times 22 \times 4 \text{ cm}$$
 Simplify
= 176 cm

4. The total surface area of a closed cylinder is given by $A = 2\pi r^2 + 2\pi rh$, where r is the radius of the circular faces, h is the height. Use the formula to calculate the area of a cylinder with a radius of 8 cm and height of 20 cm. Use 3.14 for π .

- $A = 2\pi r^{2} + 2\pi rh$ = 2 × 3.14 × 8² + 2 × 3.14 × 8 × 20 = 401.92 + 1,004.8 = 1,406.72 cm² Formula Substitute $\pi = 3.14, r = 8, h = 20$ Simplify
- 5. The volume of a cylinder is given by $V = \pi r^2 h$, where *r* is the radius of the circular base and *h* is the height. Use $\pi = \frac{22}{7}$ to find the volume of a cylinder with the base radius of 7 cm and height of 40 cm.

Solution:

V	$=\pi r^2 h$	Formula
	$=\frac{22}{7} \times 7 \text{ cm} \times 7 \text{ cm} \times 40 \text{ cm}$	Substitute $\pi = \frac{22}{7}, r = 7, h = 40$
	$= 22 \times 7 \text{ cm} \times 40 \text{ cm}$	Simplify
	$= 6,160 \text{ cm}^3$	

- 6. Given that $E = v + \frac{1}{2}mv^2$, find the value of *E* if v = 10 and m = 4. Solution:
 - *E* = $v + \frac{1}{2}mv^2$ Formula = $10 + \frac{1}{2} \times 4 \times 10^2$ Substitute v = 10, m = 4= $10 + 2 \times 100$ Simplify = 10 + 200= 210

Practice

- 1. Given that x = -1, y = 3, and z = 5 find the value of the algebraic expressions:
 - a. *y* − 3*x*
 - b. xy + z
 - c. $2y + z^2$
 - d. $x^2 5x + 2$
- 2. The relationship between the Celsius (*C*) and Fahrenheit (*F*) scales of temperature is given by $C = \frac{5}{9}(F 32)$. Find the temperature in Celsius if $F = 42^{\circ}$ F.

- 3. Given that v = u + at, find the value of v if u = 10, a = 30, $t = \frac{1}{6}$
- 4. The length of a rectangle is given by the formula $l = \frac{P}{2} w$, where *P* is the perimeter and *w* is the width. Find the value of *l* when *P* = 74 cm and *w* = 15 cm.
- 5. The formula $I = \frac{P \times R \times T}{100}$, find *I* if P = 450,000; R = 4% and T = 2.
- 6. The curved surface area of a cylinder is given by $A = 2\pi rh$, where *r* is a radius of circular base and *h* is the height. Find the area if the radius is 15 cm and height is 25 cm. Use 3.14 for π .

Lesson Title: Change of subject – Part 1	Theme: Algebraic Processes
Practice Activity: PHM1-L085	Class: SSS 1

By the end of the lesson, you will be able to change the subject of a formula.

Overview

A formula is a type of equation which shows the relationship between variables such as x and y. For an equation to be a formula, it must have more than one variable. For example, m + n = 9 is a formula, because it shows the relationship between m and n. 3n - 4 = 0 is not a formula because it only has one variable, n.

The subject of a formula is the single variable to which everything else in the formula is equal. The subject of a formula will usually be positioned to the left of the equals sign. For example, x is the subject of the formula x = 3 + y.

We can change the subject of a formula. For example, consider x = 3 + y. We could make *y* the subject. Simply subtract 3 from both sides, and we have x - 3 = y. This can also be written as y = x - 3. This is the same formula as x = 3 + y, but we have changed the subject from *x* to *y*.

We change the subject by getting a variable by itself on one side of the equals sign. We may use addition, subtraction, multiplication, or division to do this. Remember the following rules when changing the subject of an equation:

- Adding and subtracting are the opposite of one another
- Multiplying and dividing are the opposite of one another
- You must perform the same method on both sides of the equation.

If you are trying to change the subject to a variable that appears more than once in a formula, first collect all terms that contain that variable on one side of the equation. Combine like terms or factor if possible, and solve.

Solved Examples

- 1. Make *x* the subject of each formula:
 - a. y = 4x; b. y = x + 2 c. $y = \frac{1}{2}x$ d. $y = \frac{x-1}{2}$

Solutions:

a. To get *x* by itself, divide throughout by 4:

$$y = 4x$$

$$\frac{y}{4} = \frac{4x}{4}$$

$$\frac{y}{4} = x$$

b. To solve for *x*, subtract 2 from both sides:

$$y = x + 2$$

$$y - 2 = x + 2 - 2$$

$$y - 2 = x$$

c. To solve for *x*, multiply both sides by 2:

$$y = \frac{1}{2}x$$

$$2y = 2\left(\frac{1}{2}x\right)$$

$$2y = x$$

d. To write the formula with x as the subject, we must take 2 steps:

 $y = \frac{x-1}{2}$ $2y = 2\left(\frac{x-1}{2}\right)$ 2y = x-1 2y + 1 = x - 1 + 1Add 1 to both sides 2y + 1 = x

2. Make *x* the subject of the formula z = 4x + 4y. Solution:

We must take multiple steps to get *x* by itself:

z = 4x + 4y z - 4y = 4x + 4y - 4ySubtract 4y from both sides z - 4y = 4x $\frac{z - 4y}{4} = \frac{4x}{4}$ Divide both sides by 4 $\frac{z - 4y}{4} = x$

3. Make y the subject of 3y + 3x = 2y + 1. Solution:

The variable y appears twice. We must collect all terms containing y on one side of the equation before solving.

3y + 3x	=	2y + 1	
3y - 2y	=	-3x + 1	Group terms containing y
у	=	-3x + 1	Combine like terms

4. Make x the subject of ny + 3x = tx + 2. Solution:

The variable x appears twice. We must collect all terms containing x on one side of the equation before solving.

ny + 3x	=	tx + 2	
3x - tx	=	2 - ny	Group terms containing x
x(3-t)	=	2 - ny	Factorise x

$$\frac{x(3-t)}{3-t} = \frac{2-ny}{3-t}$$
$$x = \frac{2-ny}{3-t}$$

Divide both sides by 3 - t

5. Make *c* the subject of the formula $A = \frac{1}{2}(b + c)h$ Solution:

$$A = \frac{1}{2}(b+c)h$$

$$2A = 2 \times \frac{1}{2}(b+c)h$$
Multiply both sides by 2
$$2A = (b+c)h$$

$$\frac{2A}{h} = \frac{(b+c)h}{h}$$
Divide both sides by h
$$\frac{2A}{h} = b+c$$

$$\frac{2A}{h} - b = b-b+c$$
Subtract b from both sides
$$\frac{2A}{h} - b = c$$

6. Making *y* the subject of m + n + ey = 2m - ySolution:

$$m + n + ey = 2m - y$$

$$ey + y = 2m - m - n$$
 Group terms containing y

$$y(e + 1) = 2m - m - n$$
 Factorise y

$$y(e + 1) = m - n$$
 Combine like terms with m

$$\frac{y(e+1)}{e+1} = \frac{m - n}{e+1}$$
 Divide both sides by $e + 1$

$$y = \frac{m - n}{e+1}$$

Practice

1. Make *b* the subject of each formula:

a.
$$a = 13b$$
; b. $a = b - 3$ c. $a = \frac{1}{4}b$ d. $a = 2b - 8$

- 2. Make *a* the subject in the formula p = 2(a + b).
- 3. Make *y* the subject in the following relations:
 - a. 4d + 5cy = qy + rt

b.
$$3py - ay = r + s$$

- c. ky + 6x = 4y 3x
- 4. Make *u* the subject of the following:
 - a. $v = \frac{u+1}{u+2}$
b. $hu = \frac{w+3u}{m}$
c. $n = \frac{2uk}{v} + 2$
- 5. Make *F* the subject of the formula $C = \frac{5}{9}(F 32)$.

Lesson Title: Change of subject – Part 2	Theme: Algebraic Processes
Practice Activity: PHM1-L086	Class: SSS 1

By the end of the lesson, you will be able to change the subject of a formula.

Overview

This lesson is on changing the subject of more complicated relations, involving powers and roots.

You must always ensure that whatever you do in changing the subject of a formula, you maintain balance in your equation by carrying out the same operations on both sides of the equal sign. For example:

- Adding or subtracting the same quantity on both sides
- Multiplying or dividing both sides by the same quantity
- Applying the same power or root to both sides.

Note that squaring "undoes" a square root, and taking the square root "undoes" a square.

Solved Examples

1. Make *q* the subject of the relation $p = \sqrt{mq}$

Solution:

 $p = \sqrt{mq}$ $p^{2} = (\sqrt{mq})^{2}$ Square both sides to remove the square root $p^{2} = mq$ $\frac{p^{2}}{m} = \frac{mq}{m}$ Divide both sides by *m* to isolate *q* as subject $q = \frac{p^{2}}{m}$

2. Make *b* the subject of the formula $p = \sqrt{a+b}$ Solution:

 $p = \sqrt{a+b}$ $p^{2} = (\sqrt{a+b})^{2}$ Square both sides $p^{2} = a+b$ $p^{2}-a = b$

3. Make *i* the subject of the formula $R = (1 + i)^2$. $R = (1 + i)^2$
$\begin{array}{rcl} \sqrt{R} & = & \sqrt{(1+i)^2} & & \mbox{Take the square root of both sides} \\ \sqrt{R} & = & 1+i & \\ i & = & \sqrt{R}-1 & & \mbox{Subtract 1 from each side} \end{array}$

4. Make *R* the subject of the relation $q = \frac{m}{R^2} + k$ Solution: $q = \frac{m}{R^2} + k$

$$q - k = \frac{m}{R^2}$$
Subtract k from both sides

$$R^2 q - kR^2 = R^2 \left(\frac{m}{R^2}\right)$$
Multiply both sides by R^2 to remove fraction

$$R^2 (q - k) = m$$
Factorise R^2

$$\frac{R^2 (q - k)}{q - k} = \frac{m}{q - k}$$
Divide both sides by $q - k$

$$R^2 = \frac{m}{q - k}$$
Take the square root of both sides

Remember that when you find the square root of a number, the result can be either positive or negative.

5. Make *y* the subject of the relation, $N + \sqrt{y} = q$. Then, find the value of *y* when q = 5, N = 3.

Solution:

Step 1. Change the subject:

 $N + \sqrt{y} = q$ $\sqrt{y} = q - N$ Subtract *N* from both sides $\left(\sqrt{y}\right)^2 = (q - N)^2$ Square both sides $y = (q - N)^2$ Substitute the given values:

Step 2. Substitute the given values:

$$y = (5-3)^2$$
 Substitute $q = 5$ and $N = 3$
 $y = 2^2 = 4$

6. Make *x* the subject of the relation $y = ax^2 + bd + c$ Solution:

$$y = ax^{2} + bd + c$$

$$y - bd - c = ax^{2}$$
Subtract a
$$\frac{y - bd - c}{a} = \frac{ax^{2}}{a}$$
Divide bo
$$\frac{y - bd - c}{a} = x^{2}$$

$$x = \pm \sqrt{\frac{y - bd - c}{a}}$$
Take the

Subtract bd and c from both sides Divide both sides by a

Take the square root of both sides

Practice

- 1. Solve for $r: p = r^2 + 1$
- 2. Make *t* the subject of the formula $x = \sqrt{t + u}$.
- 3. Make *x* the subject of the relation $y = \frac{1}{3}\sqrt{x+1}$. Then, find the value of *x* when y = -2.
- 4. Solve for u: $F = \frac{mv^2}{u}$
- 5. Make *t* the subject of the formula $x = uf + \frac{1}{2}gt^2$.
- 6. Make *r* the subject of the relation $\frac{1}{3}\pi r^2 = v$.
- 7. Solve for *u*: $c = \sqrt{\frac{u-a}{u-b}}$

Lesson Title: Reduction to basic form of	Theme: Numbers and Numeration
surds	
Practice Activity: PHM1-L087	Class: SSS 1
Ť	

By the end of the lesson, you will be able to reduce surds to basic forms.

Overview

Surds are numbers that we cannot find a whole number square root of. They can be left in square root form to express their exact values. For example, consider these square roots: $\sqrt{2}$, $\sqrt{3}$, $\sqrt{5}$. These are examples of surds, because we cannot simplify them to whole numbers.

Surds are left in root form (with $\sqrt{}$) to express their exact value. When calculated, surds have an infinite number of non-recurring decimals. Therefore, surds are irrational numbers. The square roots of all prime numbers are surds. They give decimals that never repeat and never end.

In simplifying surds to basic forms, it is often necessary to find the largest perfect square factor. You will use the fact that $\sqrt{a \times b} = \sqrt{a} \times \sqrt{b}$. For example, consider $\sqrt{8}$. This is a surd that can be simplified. Note that 4 divides 8, and 4 is a perfect square.

Simplify $\sqrt{8}$ as follows: $\sqrt{8} = \sqrt{4 \times 2} = \sqrt{4} \times \sqrt{2} = 2\sqrt{2}$.

Always remember some common perfect squares. This will help you solve problems with surds. Some common perfect squares are listed here:

Number	Square	Number	Square	Number	Square
2	4	7	49	12	144
3	9	8	64	13	169
4	16	9	81	14	196
5	25	10	100	15	225
6	36	11	121		

Solved Examples

1. Reduce each of the following surds to its basic form:

b. $\sqrt{12}$

a. $\sqrt{8}$ Solutions:

b.	$\sqrt{12}$	$=\sqrt{4 \times 3}$
		$=\sqrt{4} \times \sqrt{3}$
		$= 2 \times \sqrt{3}$
	b.	b. √12

c. $\sqrt{18}$

d. $\sqrt{24}$

$$= 2\sqrt{2} \qquad = 2\sqrt{3}$$

c. $\sqrt{18} = \sqrt{9 \times 2}$
 $= \sqrt{9} \times \sqrt{2}$
 $= 3 \times \sqrt{2}$
 $= 3\sqrt{2}$
d. $\sqrt{24} = \sqrt{4 \times 6}$
 $= \sqrt{4} \times \sqrt{6}$
 $= 2 \times \sqrt{6}$
 $= 2\sqrt{6}$

2. Reduce each surd to its basic form:

a.
$$\sqrt{288}$$
 b. $\sqrt{192}$ c. $\sqrt{48}$ d. $\sqrt{125}$
Solutions:
a. $\sqrt{288} = \sqrt{144 \times 2}$
 $= \sqrt{144} \times \sqrt{2}$
 $= 12 \times \sqrt{2}$
 $= 12 \sqrt{2}$
b. $\sqrt{192} = \sqrt{64 \times 3}$
 $= \sqrt{64} \times \sqrt{3}$
 $= 8 \times \sqrt{3}$
 $= 8\sqrt{3}$
c. $\sqrt{48} = \sqrt{16 \times 3}$
 $= \sqrt{16} \times \sqrt{3}$
 $= 4 \times \sqrt{3}$
 $= 4\sqrt{3}$
d. $\sqrt{125} = \sqrt{25 \times 5}$
 $= 5 \times \sqrt{5}$
 $= 5\sqrt{5}$

3. Simplify: $\sqrt{16,000}$

Solution:

When you encounter a large number in a surd, try to break it down into factors as usual. You may have to try factoring it a few different ways before you find the way that gives perfect squares and allows you to simplify.

$$\sqrt{16,000} = \sqrt{1,600 \times 10}
 = \sqrt{16 \times 100 \times 10}
 = \sqrt{16} \times \sqrt{100} \times \sqrt{10}
 = 4 \times 10 \times \sqrt{10}
 = 40\sqrt{10}$$

Practice

- 1. Reduce the following surds to their basic forms:
- a. $\sqrt{32}$ b. $\sqrt{40}$ c. $\sqrt{50}$ d. $\sqrt{27}$ 2. Reduce the following surds to their basic forms: a. $\sqrt{432}$ b. $\sqrt{180}$ c. $\sqrt{1,250}$ d. $\sqrt{24}$
- a. $\sqrt{432}$ b. $\sqrt{180}$ c. $\sqrt{1,250}$ d. $\sqrt{240}$ 3. Simplify the following:
 - a. $\sqrt{6,250}$ b. $\sqrt{243}$ c. $\sqrt{9,000}$ d. $\sqrt{800}$

Lesson Title: Addition and subtraction	Theme: Numbers and Numeration
Practice Activity: PHM1-L088	Class: SSS 1

By the end of the lesson, you will be able to solve simple problems involving addition and subtraction of surds.

Overview

Surds that are in basic form can be added or subtracted. We can only add or subtract like surds. Like surds have the same surd part. To add them, keep the same surd part, and add the coefficients. For example, $m\sqrt{k} + n\sqrt{k} = (m + n)\sqrt{k}$.

Mixed surds cannot be added or subtracted. That is, they cannot be simplified further. For example, $m\sqrt{k} + n\sqrt{p}$ or $m\sqrt{k} - n\sqrt{p}$ cannot be simplified further.

To add or subtract two or more surds, follow the steps below:

- Step 1. Convert each surd to its simplest form.
- Step 2. Add or subtract any like surds.

Solved Examples

1. Simplify: $3\sqrt{2} + 7\sqrt{2}$

Solution:

These are like surds, so the coefficients can be added:

$$3\sqrt{2} + 7\sqrt{2} = (3+7)\sqrt{2}$$

= $10\sqrt{2}$

2. Simplify: $\sqrt{8} + \sqrt{32}$ Solution:

$$\sqrt{8} + \sqrt{32} = \sqrt{4 \times 2} + \sqrt{16 \times 2}$$

Simplify each surd
$$= \sqrt{4} \times \sqrt{2} + \sqrt{16} \times \sqrt{2}$$

$$= 2\sqrt{2} + 4\sqrt{2}$$

$$= (2 + 4)\sqrt{2}$$

$$= 6\sqrt{2}$$

Add like surds

3. Simplify: $2\sqrt{3} - \sqrt{3}$ Solution:

These are like surds, so the coefficients can be subtracted:

$$2\sqrt{3} - \sqrt{3} = (2 - 1)\sqrt{3}$$

= $1\sqrt{3}$

 $= \sqrt{3}$ 4. Simplify: $\sqrt{27} - \sqrt{12}$ Solution: $\sqrt{27} - \sqrt{12} = \sqrt{9 \times 3} - \sqrt{4 \times 3}$ Simplify each surd $= \sqrt{9} \times \sqrt{3} - \sqrt{4} \times \sqrt{3}$ $= 3\sqrt{3} - 2\sqrt{3}$ $= (3 - 2)\sqrt{3}$ $= \sqrt{3}$

Simplify:
$$\sqrt{24} - 3\sqrt{6}$$

Solution:
 $\sqrt{24} - 3\sqrt{6} = \sqrt{4 \times 6} - 3\sqrt{6}$ Simplify $\sqrt{24}$
 $= \sqrt{4} \times \sqrt{6} - 3\sqrt{6}$
 $= 2\sqrt{6} - 3\sqrt{6}$
 $= (2 - 3)\sqrt{6}$ Subtract like surds
 $= -1\sqrt{6}$
 $= -\sqrt{6}$

Simplify
$$\sqrt{98} - 2\sqrt{8}$$

Solution:
 $\sqrt{98} - 2\sqrt{8} = \sqrt{49 \times 2} - 2\sqrt{4 \times 2}$ Simplify $\sqrt{24}$
 $= \sqrt{49} \times \sqrt{2} - 2\sqrt{4} \times \sqrt{2}$
 $= 7\sqrt{2} - 2(2)\sqrt{2}$
 $= 7\sqrt{2} - 4\sqrt{2}$
 $= (7 - 4)\sqrt{2}$ Subtract like surds
 $= 3\sqrt{2}$

Practice

5.

6.

Simplify the following:

1.
$$5\sqrt{2} - \sqrt{2}$$

2. $4\sqrt{5} + 3\sqrt{5}$
3. $\sqrt{3} + \sqrt{3}$
4. $\sqrt{24} - 2\sqrt{6}$
5. $\sqrt{50} - \sqrt{8}$
6. $\sqrt{75} - 2\sqrt{12}$
7. $3\sqrt{8} + \sqrt{50}$

Lesson Title: Addition and subtraction	Theme: Numbers and Numeration
of surds – Part 2	
Practice Activity: PHM1-L089	Class: SSS 1

By the end of the lesson, you will be able to solve more complicated problems involving addition and subtraction of surds.

Overview

Remember that we can only add or subtract surds which are like, or have the same surd part. First simplify surds to their basic form, then add or subtract if possible.

Solved Examples

1. Simplify
$$\sqrt{147} - \sqrt{75} + \sqrt{27}$$

Solution:
 $\sqrt{147} - \sqrt{75} + \sqrt{27} = \sqrt{49 \times 3} - \sqrt{25 \times 3} + \sqrt{9 \times 3}$ Simplify surds
 $= \sqrt{49 \times \sqrt{3} - \sqrt{25} \times \sqrt{3} + \sqrt{9 \times \sqrt{3}}}$
 $= 7\sqrt{3} - 5\sqrt{3} + 3\sqrt{3}$
 $= 7\sqrt{3} - 5\sqrt{3} + 3\sqrt{3}$
 $= (7 - 5 + 3)\sqrt{3}$ Add/subtract coefficients
 $= 5\sqrt{3}$
2. Simplify $\sqrt{180} - \sqrt{45} + \sqrt{20}$
Solution:
 $\sqrt{180} - \sqrt{45} + \sqrt{20} = \sqrt{36 \times 5} - \sqrt{9 \times 5} + \sqrt{4 \times 5}$ Simplify surds
 $= \sqrt{36} \times \sqrt{5} - \sqrt{9} \times \sqrt{5} + \sqrt{4} \times \sqrt{5}$
 $= 6\sqrt{5} - 3\sqrt{5} + 2\sqrt{5}$
 $= (6 - 3 + 2)\sqrt{3}$ Add/subtract
 $= 5\sqrt{5}$
3. Simplify $\sqrt{50} - 3\sqrt{2} + \frac{1}{2}\sqrt{8}$
Solution:
 $\sqrt{50} - 3\sqrt{2} + \frac{1}{2}\sqrt{8} = \sqrt{25 \times 2} - 3\sqrt{2} + \frac{1}{2}\sqrt{4 \times 2}$ Simplify surds
 $= 5\sqrt{2} - 3\sqrt{2} + \frac{2}{2}\sqrt{2}$
 $= 5\sqrt{2} - 3\sqrt{2} + \sqrt{2}$
 $= (5 - 3 + 1)\sqrt{2}$ Add/subtract
 $= 3\sqrt{2}$
4. Simplify $\frac{5}{2}\sqrt{32} - \frac{1}{5}\sqrt{125} - \frac{1}{3}\sqrt{72} + \frac{3}{2}\sqrt{180}$
Solution:

$$\frac{5}{2}\sqrt{32} - \frac{1}{5}\sqrt{125} - \frac{1}{3}\sqrt{72} + \frac{3}{2}\sqrt{180} = \frac{5}{2}\sqrt{16 \times 2} - \frac{1}{5}\sqrt{25 \times 5} - \frac{1}{3}\sqrt{36 \times 2} + \frac{3}{2}\sqrt{36 \times 5}$$
$$= \frac{5}{2} \times 4\sqrt{2} - \frac{1}{5} \times 5\sqrt{5} - \frac{1}{3} \times 6\sqrt{2} + \frac{3}{2} \times 6\sqrt{5}$$
$$= 10\sqrt{2} - \sqrt{5} - 2\sqrt{2} + 9\sqrt{5}$$
$$= 10\sqrt{2} - 2\sqrt{2} + 9\sqrt{5} - \sqrt{5}$$
$$= (10 - 2)\sqrt{2} + (9 - 1)\sqrt{5}$$
$$= 8\sqrt{2} + 8\sqrt{5}$$
$$= 8(\sqrt{2} + \sqrt{5})$$

5. If $\sqrt{294} + \sqrt{216} - \sqrt{486} = m\sqrt{6}$. Find the value of *m*. Solution:

Simplify the left-hand side, then add and subtract the surds. Based on the righthand side, it looks like we will have $\sqrt{6}$ in the surds on the left-hand side.

$$\sqrt{294} + \sqrt{216} = \sqrt{49 \times 6} + \sqrt{36 \times 6} - \sqrt{81 \times 6}$$
$$-\sqrt{486} = \sqrt{49} \times \sqrt{6} + \sqrt{36} \times \sqrt{6} - \sqrt{81} \times \sqrt{6}$$
$$= 7\sqrt{6} + 6\sqrt{6} - 9\sqrt{6}$$
$$= (7 + 6 - 9)\sqrt{6} = 4\sqrt{6}$$

Hence m = 4

Practice

Simplify the following surds as far as possible.

1.
$$\frac{2}{3}\sqrt{63} + \frac{1}{2}\sqrt{28} - \frac{1}{10}\sqrt{175}$$

2. $\frac{5}{4}\sqrt{32} - \frac{1}{3}\sqrt{72}$
3. $2\sqrt{18} - \frac{1}{2}\sqrt{128} + \frac{3}{2}\sqrt{72}$
4. $\frac{5}{2}\sqrt{48} + \frac{1}{3}\sqrt{216} - \frac{2}{5}\sqrt{75} - \frac{1}{2}\sqrt{24}$
5. $5\sqrt{125} - 2\sqrt{500} + 2\sqrt{20}$
6. $3\sqrt{7} - \sqrt{343} + 4\sqrt{28}$
7. If $\sqrt{175} - \sqrt{448} + \sqrt{112} = n\sqrt{7}$, find the value of n .

Lesson Title: Properties of surds	Theme: Numbers and Numeration
Practice Activity: PHM1-L090	Class: SSS 1

By the end of the lesson, you will be able to identify properties of surds.

Overview

Recall that surds are square roots of numbers that are imperfect squares. They are irrational numbers. The following are examples of surds: $\sqrt{2}$, $\sqrt{3}$, $\sqrt{5}$, $\sqrt{6}$, $\sqrt{7}$

In operating with surds, please note carefully the following properties of surds:

4.
$$a\sqrt{c} \pm b\sqrt{c} = (a \pm b)\sqrt{c}$$

5.
$$\sqrt{a^2} = \sqrt{a} \times \sqrt{a} = a$$

$$6. \quad a\sqrt{b} \times \sqrt{c} = a\sqrt{bc}$$

7.
$$a \times \sqrt{b} = a\sqrt{b}$$

8.
$$a \times b\sqrt{c} = ab\sqrt{c}$$

Solved Examples

1. Simplify $\sqrt{27}$ Solution:

$$\sqrt{27} = \sqrt{9 \times 3}$$

= $\sqrt{9} \times \sqrt{3}$ Using 1) $\sqrt{a \times b} = \sqrt{a} \times \sqrt{b}$
= $3 \times \sqrt{3}$
= $3\sqrt{3}$ Using 7) $a \times \sqrt{b} = a\sqrt{b}$

2. Simplify $\sqrt{3\frac{3}{8}}$

Solution:

First, change the mixed fraction to an improper fraction so that property 2. can be applied.

$$\sqrt{3\frac{3}{4}} = \sqrt{\frac{15}{4}}$$
$$= \frac{\sqrt{15}}{\sqrt{4}}$$
 Using 2) $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$

$$=\frac{\sqrt{15}}{2}$$

3. Simplify $5\sqrt{5} + 4\sqrt{5}$ Solution: $5\sqrt{5} + 4\sqrt{5} = (5+4)\sqrt{5}$ $= 9\sqrt{5}$

Using 4)
$$a\sqrt{c} + b\sqrt{c} = (a+b)\sqrt{c}$$

4. Simplify:

a.
$$3 \times 2\sqrt{5}$$
 c. $\sqrt{7^2}$
b. $2\sqrt{5} \times \sqrt{2}$ d. $\sqrt{x^3}$
Solutions:
a. $3 \times 2\sqrt{5} = (3 \times 2)\sqrt{2}$ Using 8) $a \times b\sqrt{c} = ab\sqrt{c}$
 $= 6\sqrt{2}$
b. $2\sqrt{5} \times \sqrt{2} = 2\sqrt{5 \times 2}$ Using 6) $a\sqrt{b} \times \sqrt{c} = a\sqrt{bc}$
 $= 2\sqrt{10}$
c. $\sqrt{7^2} = \sqrt{7} \times \sqrt{7}$ Using 5) $\sqrt{a^2} = \sqrt{a} \times \sqrt{a} = a$
 $= 7$
d. $\sqrt{x^3} = \sqrt{x} \times \sqrt{x} \times \sqrt{x}$ Using 5) $\sqrt{a^2} = \sqrt{a} \times \sqrt{a} = a$
 $= x \times \sqrt{x}$
 $= x\sqrt{x}$

Practice

1. Simplify: a.
$$\sqrt{32}$$
 b. $\sqrt{48}$ c. $\sqrt{24}$
2. Simplify: a. $\sqrt{2\frac{1}{4}}$ b. $\sqrt{5\frac{5}{9}}$ c. $\sqrt{1\frac{9}{16}}$
3. Simplify: a. $a\sqrt{6} + 2a\sqrt{6} - 4a\sqrt{6}$
b. $5\sqrt{2} + 6\sqrt{2} - 10\sqrt{2}$
c. $\sqrt{12} + \sqrt{27} - \sqrt{48}$
4. Simplify: a. $4 \times 2\sqrt{2}$ b. $4\sqrt{3} \times 2\sqrt{2}$ c. $\sqrt{a^5}$ d. $\sqrt{x^4}$

Lesson Title: Multiplication of surds – Part 1	Theme: Numbers and Numeration
Practice Activity: PHM1-L091	Class: SSS 1

By the end of the lesson, you will be able to multiply surds.

Overview

In the multiplication of surds you need to carefully follow the following steps:

- 1. Express each surd in its basic or simplest form.
- 2. Multiply the coefficients, and multiply the surd parts.
- 3. Make sure the result is in its simplest form, and simplify it if possible.

The general rules for multiplying surds are:

$$\sqrt{a} \times \sqrt{b} = \sqrt{ab}$$
$$a\sqrt{b} \times c\sqrt{d} = ac\sqrt{bd}$$

Solved Examples

1. Find the product of $7\sqrt{2}$ and $5\sqrt{3}$. **Solution:**

 $7\sqrt{2} \times 5\sqrt{3} = (7 \times 5)\sqrt{2 \times 3}$ Multiply coefficients and numbers in surds = $35\sqrt{6}$

The result is already in its simplest form, so the answer is $35\sqrt{6}$.

2. Simplify: $\sqrt{5} \times \sqrt{15}$

Solution:

 $\sqrt{5} \times \sqrt{15} = \sqrt{5 \times 15}$ Multiply numbers in surds $= \sqrt{75}$ $= \sqrt{25 \times 3}$ Simplify $\sqrt{75}$ $= \sqrt{25} \times \sqrt{3}$ $= 5\sqrt{3}$

3. Find the product of $4\sqrt{7}$ and $3\sqrt{5}$. **Solution:**

 $4\sqrt{7} \times 3\sqrt{5} = (4 \times 3)\sqrt{7 \times 5}$ Multiply coefficients and numbers in = $12\sqrt{35}$ surds

4. Simplify: $\sqrt{3} \times \sqrt{18}$ Solution: $\sqrt{3} \times \sqrt{18} = \sqrt{3} \times \sqrt{9 \times 2}$ Simplify $\sqrt{18}$

$$= \sqrt{3} \times (\sqrt{9} \times \sqrt{2})$$

= $\sqrt{3} \times 3\sqrt{2}$
= $3\sqrt{3} \times 2$
= $3\sqrt{6}$
Multiply numbers in surds

5. Multiply:
$$\sqrt{6} \times \sqrt{18}$$

Solution:

$$\sqrt{6} \times \sqrt{18} = \sqrt{6} \times \sqrt{9 \times 2}$$

Simplify $\sqrt{18}$
$$= \sqrt{6} \times (\sqrt{9} \times \sqrt{2})$$

$$= \sqrt{6} \times 3\sqrt{2}$$

$$= 3\sqrt{6 \times 2}$$

Multiply numbers in surds
$$= 3\sqrt{12}$$

$$= 3\sqrt{4 \times 3}$$

Simplify $3\sqrt{12}$
$$= 3\sqrt{4}\sqrt{3}$$

$$= 3 \times 2\sqrt{3}$$

$$= 6\sqrt{3}$$

6. Multiply: $3\sqrt{24} \times 2\sqrt{12}$ Solution:

$$3\sqrt{24} \times = 3\sqrt{4 \times 6} \times 2\sqrt{4 \times 3}$$

Simplify both surds
$$= 3\sqrt{4} \times \sqrt{6} \times 2 \times \sqrt{4} \times \sqrt{3}$$

$$= 3 \times 2 \times \sqrt{6} \times 2 \times 2 \times \sqrt{3}$$

$$= 6\sqrt{6} \times 4\sqrt{3}$$

$$= 6 \times 4\sqrt{6 \times 3}$$

Multiply
$$= 24\sqrt{18}$$

$$= 24 \times \sqrt{9 \times 2}$$

Simplify
$$= 24 \times 3\sqrt{2}$$

$$= 72\sqrt{2}$$

7. Simplify: $(2\sqrt{3})^2$ Solution:

$$(2\sqrt{3})^2 = 2\sqrt{3} \times 2\sqrt{3}$$
 Multiply
= $(2 \times 2)\sqrt{3 \times 3}$ Multiply coefficients and numbers in surds
= $4\sqrt{9}$
= 4×3
= 12

Practice

- 1. Find the product of:
 - a. $7\sqrt{3}$ and $5\sqrt{3}$
 - b. $\sqrt{32}$ and $\sqrt{80}$
- 2. Simplify:
 - a. $4\sqrt{2} \times 2\sqrt{8}$
 - b. $4\sqrt{12} \times 6\sqrt{18}$
- 3. Multiply:
 - a. $\sqrt{64} \times \sqrt{48}$
 - b. $2\sqrt{75} \times \sqrt{125}$
- 4. Simplify:

a.
$$(3\sqrt{2})^2$$

b. $\sqrt{54} \times \sqrt{128}$

Lesson Title: Multiplication of surds – Part 2	Theme: Numbers and Numeration
Practice Activity: PHM1-L092	Class: SSS 1

By the end of the lesson, you will be able to multiply surds.

Overview

The lesson today is on the simplification of more complicated problems involving the multiplication of surds.

Remember the basic laws involving multiplication of surds. The coefficient parts are multiplied, and the surd parts are multiplied. For example:

a.
$$\sqrt{a} \times \sqrt{b} = \sqrt{a \times b} = \sqrt{ab}$$

b. $a \times b\sqrt{c} = ab\sqrt{c}$
c. $a\sqrt{b} \times c\sqrt{d} = ab\sqrt{bd}$

Remember the order of operations (BODMAS) and apply it when there are multiple operations in one problem.

Solved Examples

1. Simplify:

a.
$$\sqrt{24} \times (\sqrt{3})^2$$

b. $10(\sqrt{3} + 2)$
Solution:

a.
$$\sqrt{24} \times (\sqrt{3})^2 = \sqrt{4 \times 6} \times (\sqrt{3} \times \sqrt{3})$$

 $= \sqrt{4} \times \sqrt{6} \times 3$
 $= 2 \times 3 \times \sqrt{6}$
 $= 6\sqrt{6}$
b. $10(\sqrt{3}+2) = 10\sqrt{3} + 10(2)$
 $= 10\sqrt{3} + 20$
 $= 20 + 10\sqrt{3}$

2. Simplify: $\sqrt{10}(\sqrt{8} + 2)$ Solution:

$$\sqrt{10}(\sqrt{8} + 2) = \sqrt{10} \times \sqrt{8} + 2 \times \sqrt{10}$$
$$= \sqrt{10 \times 8} + 2\sqrt{10}$$
$$= \sqrt{80} + 2\sqrt{10}$$
$$= \sqrt{16 \times 5} + 2\sqrt{10}$$
$$= \sqrt{16} \times \sqrt{5} + 2\sqrt{10}$$
$$= 4\sqrt{5} + 2\sqrt{10}$$

3. Simplify: $\sqrt{48} - 3\sqrt{2}(2\sqrt{3} - 4) - 4\sqrt{48}$ Solution:

$$\begin{array}{rcl} \sqrt{48} - 3\sqrt{2}(2\sqrt{3} - 4) - 4\sqrt{48} &= \sqrt{16 \times 3} - 3 \times 2\sqrt{2 \times 3} + 3 \times 4\sqrt{2} - 4\sqrt{16 \times 3} \\ &= \sqrt{16} \times \sqrt{3} - 6\sqrt{6} + 12\sqrt{2} - 4 \times \sqrt{16} \times \sqrt{3} \\ &= 4\sqrt{3} - 6\sqrt{6} + 12\sqrt{2} - 4 \times 4\sqrt{3} \\ &= 4\sqrt{3} - 6\sqrt{6} - 12\sqrt{2} - 16\sqrt{3} \\ &= -12\sqrt{2} - 12\sqrt{3} - 6\sqrt{6} \end{array}$$

4. Simplify $(\sqrt{2} - 1)(3 - \sqrt{2})$ and write the result in the form $a + b\sqrt{c}$, stating the value of *a*, *b* and *c*. **Solution:**

$$(\sqrt{2} - 1)(3 - \sqrt{2}) = \sqrt{2}(3 - \sqrt{2}) - 1(3 - \sqrt{2})$$

= $3\sqrt{2} - \sqrt{2} \times \sqrt{2} - 3 + \sqrt{2}$
= $3\sqrt{2} - 2 - 3 + \sqrt{2}$
= $-5 + 4\sqrt{2}$

Hence a = -5, b = 4 and c = 2

5. Simplify: $\frac{\sqrt{125} \times \sqrt{63}}{\sqrt{28} \times \sqrt{80}}$.

Solution:

$$\frac{\sqrt{125} \times \sqrt{63}}{\sqrt{28} \times \sqrt{80}} = \frac{\sqrt{25 \times 5} \times \sqrt{9 \times 7}}{\sqrt{4 \times 7} \times \sqrt{16 \times 5}}$$
$$= \frac{\sqrt{25} \times \sqrt{5} \times \sqrt{9} \times \sqrt{7}}{\sqrt{4} \times \sqrt{7} \times \sqrt{16} \times \sqrt{5}}$$
$$= \frac{5 \sqrt{5} \times 3 \sqrt{7}}{2 \sqrt{7} \times 4 \sqrt{5}}$$
$$= \frac{5 \times 3 \sqrt{7} \sqrt{5}}{2 \times 4 \sqrt{5} \sqrt{7}}$$
$$= \frac{15 \sqrt{35}}{8 \sqrt{35}}$$
$$= \frac{15}{8} = 1\frac{7}{8}$$

6. Simplify $(2 - 3\sqrt{3})(3 + 2\sqrt{3})$ and write the result in the form $a + b\sqrt{c}$. **Solution:**

Expand the brackets, then simplify:

$$(2 - 3\sqrt{3}) (3 + 2\sqrt{3}) = 2(3 + 2\sqrt{3}) - 3\sqrt{3} (3 + 2\sqrt{3})$$
$$= 6 + 4\sqrt{3} - 9\sqrt{3} - 6(3)$$
$$= (6 - 18) + (4 - 9)\sqrt{3}$$
$$= -12 - 5\sqrt{3}$$

Practice

- 1. Simplify: $\sqrt{20} \times (\sqrt{5})^2$
- 2. Simplify: $\sqrt{2}(3 2\sqrt{2})$
- 3. Simplify: $\sqrt{18} \left(\sqrt{27} \frac{2}{\sqrt{2}} \right)$
- 4. Simplify: $2\sqrt{5}(6 3\sqrt{5})$
- 5. Simplify $\sqrt{18}(3+2\sqrt{8}) + 3\sqrt{2}$ and write your result in the form $a + b\sqrt{c}$.
- 6. Simplify: $\frac{\sqrt{80} \times \sqrt{63}}{\sqrt{125} \times \sqrt{28}}$
- 7. Simplify $(2 + 3\sqrt{3})^2$ and write the result in the form $a + b\sqrt{c}$

Lesson Title: Rationalisation of denominators of surds – Part 1	Theme: Numbers and Numeration
Practice Activity: PHM1-L093	Class: SSS 1

By the end of the lesson, you will be able to rationalise the denominators of surds.

Overview

Rationalising the denominator of a surd is the process of changing the denominator from a surd to a rational number.

The denominator of a fraction should not be an irrational number. To make the irrational number in the denominator a rational number, we multiply both the numerator and denominator by the denominator.

The general rule is $\frac{a}{\sqrt{b}} = \frac{a}{\sqrt{b}} \times \frac{\sqrt{b}}{\sqrt{b}} = \frac{a\sqrt{b}}{b}$.

Remember that when you multiply a fraction that has the same numerator and denominator, you are actually multiplying by 1. We can have irrational numbers in the numerator of a fraction.

Solved Examples

1. Simplify: $\frac{1}{\sqrt{2}}$

Solution:

$$\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}}$$
 Multiply the top and bottom by $\sqrt{2}$
$$= \frac{1 \times \sqrt{2}}{\sqrt{2} \times \sqrt{2}}$$
$$= \frac{\sqrt{2}}{2}$$
 Note: $\sqrt{2} \times \sqrt{2} = 2$

2. Simplify the following:

a. $\frac{\sqrt{3}}{\sqrt{7}}$ b. $\frac{10}{\sqrt{5}}$ Solutions: a. $\frac{\sqrt{3}}{\sqrt{7}} = \frac{\sqrt{3}}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}}$ b. $\frac{10}{\sqrt{5}} = \frac{10}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$ $= \frac{\sqrt{21}}{7}$ $= \frac{10\sqrt{5}}{5}$ $= 2\sqrt{5}$

- 3. Simplify the following:
 - a. $\frac{4}{\sqrt{24}}$ b. $\frac{2\sqrt{2}}{\sqrt{12}}$ c. $\frac{3}{\sqrt{27}}$ d. $\frac{9}{\sqrt{27}}$

Solutions:

a.
$$\frac{4}{\sqrt{24}} = \frac{4}{2\sqrt{6}}$$

 $= \frac{2}{\sqrt{6}}$
 $= \frac{2}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}}$
 $= \frac{2}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}}$
 $= \frac{2\sqrt{6}}{\sqrt{6}}$
 $= \frac{\sqrt{2}}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{\sqrt{6}}{3}$
c. $\frac{3}{\sqrt{27}} = \frac{3}{3\sqrt{3}}$
 $= \frac{1}{\sqrt{3}}$
 $= \frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{\sqrt{3}}{3}$
 $= \sqrt{3}$
b. $\frac{2\sqrt{2}}{\sqrt{12}} = \frac{2\sqrt{2}}{2\sqrt{3}}$
 $= \frac{\sqrt{2}}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{\sqrt{6}}{3}$
d. $\frac{9}{\sqrt{27}} = \frac{9}{3\sqrt{3}}$
 $= \frac{3}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{3\sqrt{3}}{3}$
 $= \sqrt{3}$

4. Simplify the following:

a.
$$\frac{3\sqrt{3}}{2\sqrt{5}}$$
 b. $\frac{5}{3\sqrt{5}}$ c. $\frac{3\sqrt{3}}{2\sqrt{5}}$
Solutions:
a. $\frac{3\sqrt{3}}{2\sqrt{5}} = \frac{3\sqrt{3}}{2\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$ b. $\frac{5}{3\sqrt{5}} = \frac{5}{3\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$
 $= \frac{3\sqrt{15}}{2\times5}$ $= \frac{5\sqrt{5}}{3\times5}$
 $= \frac{3\sqrt{15}}{10}$ $= \frac{\sqrt{5}}{3}$
c. $\frac{3\sqrt{3}}{2\sqrt{5}} = \frac{3\sqrt{3}}{2\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$
 $= \frac{3\sqrt{15}}{2\times5}$
 $= \frac{3\sqrt{15}}{2\times5}$
 $= \frac{3\sqrt{15}}{10}$

Practice

Simplify the following surds:

1.
$$\frac{3}{\sqrt{8}}$$
 2. $\frac{\sqrt{5}}{\sqrt{7}}$ 3. $\frac{16}{\sqrt{128}}$ 4. $\frac{\sqrt{27}}{\sqrt{72}}$ 5. $\frac{3\sqrt{2}}{\sqrt{27}}$ 6. $\frac{2\sqrt{5}}{3\sqrt{2}}$
7. $\frac{24}{\sqrt{24}}$ 8. $\frac{7\sqrt{2}}{\sqrt{14}}$ 9. $\frac{18}{\sqrt{162}}$ 10. $\frac{4\sqrt{3}}{3\sqrt{2}}$

Lesson Title: Rationalisation of the	Theme: Numbers and Numeration
denominator of surds – Part 2	
Practice Activity: PHM1-L094	Class: SSS 1

By the end of the lesson, you will be able to rationalise the denominator of surds.

Overview

This is the second lesson on rationalising the denominator of surds. It covers problems where there are 2 terms in the numerator or denominator, and at least one of them is a surd.

We use a special process when there are 2 terms in the denominator and one of them is a surd. For example, consider $\frac{1}{3-\sqrt{2}}$. Such problems require you to multiply the numerator and denominator by the **conjugate** of the binomial. To find the conjugate, change the sign in the middle. The conjugate of $3 - \sqrt{2}$ is $3 + \sqrt{2}$.

When a binomial is multiplied by its conjugate, simply square each term to save time. For example, $(3 - \sqrt{2})(3 + \sqrt{2}) = 3^2 - \sqrt{2}^2$. Observe why this is true. The middle terms will always cancel:

$$(3 - \sqrt{2})(3 + \sqrt{2}) = 3(3 + \sqrt{2}) - \sqrt{2}(3 + \sqrt{2})$$

Multiply
$$= 3^2 + 3\sqrt{2} - 3\sqrt{2} - \sqrt{2}^2$$

Remove brackets
$$= 3^2 - \sqrt{2}^2$$

Middle terms cancel

You will also see problems with 2 terms in the numerator, such as $\frac{\sqrt{3}+\sqrt{7}}{\sqrt{21}}$. These problems are solved using the process from the previous class. Multiply the numerator and denominator by $\sqrt{21}$ to rationalise the denominator, then simplify.

Solved Examples

1. Simplify:
$$\frac{1}{3-\sqrt{2}}$$

Solution:

Note that you will need to multiply the numerator and denominator by $3 + \sqrt{2}$, the conjugate of $3 - \sqrt{2}$.

$$\frac{1}{3-\sqrt{2}} = \frac{1}{3-\sqrt{2}} \times \frac{3+\sqrt{2}}{3+\sqrt{2}}$$
$$= \frac{3+\sqrt{2}}{3^2-\sqrt{2}^2}$$
$$= \frac{3+\sqrt{2}}{9-2}$$
$$= \frac{3+\sqrt{2}}{7}$$

Multiply top and bottom by $3 + \sqrt{2}$

Simplify

2. Simplify: $\frac{3}{2+\sqrt{5}}$

Solution:

$$\frac{3}{2+\sqrt{5}} = \frac{3}{2+\sqrt{5}} \times \frac{2-\sqrt{5}}{2-\sqrt{5}} \\ = \frac{3(2-\sqrt{5})}{(2)^2 - (\sqrt{5})^2} \\ = \frac{6-3\sqrt{5}}{4-5} \\ = \frac{6-3\sqrt{5}}{-1} \\ = -(6-3\sqrt{5}) \\ = -6+3\sqrt{5}$$

3. Simplify the following:

a.
$$\frac{2}{3+2\sqrt{2}}$$
 b. $\frac{5}{9-3\sqrt{5}}$
Solutions:
a. $\frac{2}{3+2\sqrt{2}} = \frac{2}{3+2\sqrt{2}} \times \frac{3-2\sqrt{2}}{3-2\sqrt{2}}$
 $= \frac{2(3-2\sqrt{2})}{(3)^2 - (2\sqrt{2})^2}$
 $= \frac{6-4\sqrt{2}}{9-8}$
 $= \frac{6-4\sqrt{2}}{1}$
 $= 6 - 4\sqrt{2}$
b. $\frac{5}{9-3\sqrt{5}} = \frac{5}{9-3\sqrt{5}} \times \frac{9+3\sqrt{5}}{9+3\sqrt{5}}$
 $= \frac{5(9+3\sqrt{5})}{9^2 - (3\sqrt{5})^2}$
 $= \frac{45+15\sqrt{5}}{81-45}$
 $= \frac{45+15\sqrt{5}}{36}$
 $= \frac{45}{36} + \frac{15}{36}\sqrt{5}$
 $= \frac{5}{4} + \frac{5}{12}\sqrt{5}$

4. Express $\frac{\sqrt{3}+\sqrt{7}}{\sqrt{21}}$ in the form $a\sqrt{3} + b\sqrt{7}$, where *a* and *b* are rational numbers. **Solution:**

$$\frac{\sqrt{3}+\sqrt{7}}{\sqrt{21}} = \frac{\sqrt{3}+\sqrt{7}}{\sqrt{21}} \times \frac{\sqrt{21}}{\sqrt{21}}$$
Rationalise the denominator
$$= \frac{\sqrt{21}(\sqrt{3}+\sqrt{7})}{\sqrt{21}\times\sqrt{21}}$$

$$= \frac{\sqrt{63}+\sqrt{147}}{21}$$
Multiply the numerator
$$= \frac{\sqrt{9\times7}+\sqrt{49\times3}}{21}$$
Simplify surds

$$= \frac{3\sqrt{7} + 7\sqrt{3}}{21}$$
$$= \frac{7\sqrt{3} + 3\sqrt{7}}{21}$$
$$= \frac{7\sqrt{3}}{21} + \frac{3\sqrt{7}}{21}$$
$$= \frac{1}{3}\sqrt{3} + \frac{1}{7}\sqrt{7}$$

Change the order of the numerator Simplify

where
$$a = \frac{1}{3}$$
 and $b = \frac{1}{7}$

5. Simplify: $\frac{\sqrt{3}+\sqrt{5}}{\sqrt{15}}$

Solution:

$$\frac{\sqrt{3}+\sqrt{5}}{\sqrt{15}} = \frac{\sqrt{3}+\sqrt{5}}{\sqrt{15}} \times \frac{\sqrt{15}}{\sqrt{15}}$$
$$= \frac{\sqrt{15}(\sqrt{3}+\sqrt{5})}{\sqrt{15}\times\sqrt{15}}$$
$$= \frac{\sqrt{45}+\sqrt{75}}{15}$$
$$= \frac{\sqrt{9}\times5+\sqrt{25\times3}}{15}$$
$$= \frac{3\sqrt{5}+\sqrt{25\times3}}{15}$$
$$= \frac{3\sqrt{5}}{15} + \frac{5\sqrt{3}}{15}$$
$$= \frac{\sqrt{5}}{5} + \frac{\sqrt{3}}{3}$$
$$= \frac{1}{5}\sqrt{5} + \frac{1}{3}\sqrt{3}$$

6. Simplify
$$\frac{1+\sqrt{2}}{\sqrt{2}}$$
 to the form $m + n\sqrt{2}$.
Solution:

$$\begin{array}{rcl} \frac{1+\sqrt{2}}{\sqrt{2}} &= \frac{1+\sqrt{2}}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} \\ &= \frac{\sqrt{2}(1+\sqrt{2})}{\sqrt{2}\times\sqrt{2}} \\ &= \frac{\sqrt{2}+2}{2} \\ &= \frac{\sqrt{2}}{2} + \frac{2}{2} \\ &= 1 + \frac{1}{2}\sqrt{2} \ \text{ where } m = 1 \ \text{and } n = \frac{1}{2} \end{array}$$

Practice

1. Simplify:

a.
$$\frac{3}{4-\sqrt{3}}$$
 b. $\frac{1}{3+\sqrt{2}}$ c. $\frac{5}{5+\sqrt{2}}$ d. $\frac{6}{7-2\sqrt{2}}$ e. $\frac{7}{8-2\sqrt{3}}$
2. Simplify the following:
a. $\frac{9-\sqrt{108}}{\sqrt{72}}$ b. $\frac{\sqrt{5}+\sqrt{75}}{\sqrt{27}}$ c. $\frac{8+\sqrt{12}}{2\sqrt{5}}$
3. Simplify: $\frac{\sqrt{3}+\sqrt{2}}{\sqrt{24}}$

Lesson Title: Expansion and simplification of surds	Theme: Numbers and Numeration
Practice Activity: PHM1-L095	Class: SSS 1

By the end of the lesson, you will be able to expand and simplify expressions involving surds.

Overview

In expanding a bracket that involve surds you use the distributive law. For example, consider $\sqrt{5} \left(\sqrt{45} - \frac{8}{\sqrt{80}} \right)$. The term in front of the bracket ($\sqrt{5}$) will be multiplied by each term inside the brackets.

You will also see problems that involve the multiplication of 2 binomials, such as $(1 + \sqrt{5})(1 - \sqrt{5})$. Distribute as usual. Multiply each term in the first bracket by the expression in the second bracket.

Remember to follow the rules for multiplying surds. Rational numbers can be multiplied by each other, and surds can be multiplied by each other. However, numbers and surds behave like numbers and variables in algebra and cannot be combined (For example, $2 \times \sqrt{5} = 2\sqrt{5}$, and this cannot be simplified further).

Solved Examples

1. Expand and simplify
$$(1 + \sqrt{5})(1 - \sqrt{5})$$

Solution:
 $(1 + \sqrt{5})(1 - \sqrt{5}) = 1(1 - \sqrt{5}) + \sqrt{5}(1 - \sqrt{5})$ Expand (distribute)
 $= 1 - \sqrt{5} + \sqrt{5} - (\sqrt{5})^2$ Simplify
 $= 1 - \sqrt{5} + \sqrt{5} - 5$ Note that $(\sqrt{5})^2 = \sqrt{5} \times \sqrt{5} = 5$
 $= 1 - 5 - \sqrt{5} + \sqrt{5}$ Collect like terms
 $= -4$

2. Expand and simplify $(\sqrt{5} + \sqrt{2})(\sqrt{5} + \sqrt{2})$ Solution:

$$(\sqrt{5} + \sqrt{2})(\sqrt{5} + \sqrt{2}) = \sqrt{5}(\sqrt{5} + \sqrt{2}) + \sqrt{2}(\sqrt{5} + \sqrt{2})$$

$$= (\sqrt{5})^2 + \sqrt{5 \times 2} + \sqrt{2 \times 5} + (\sqrt{2})^2$$

$$= 5 + \sqrt{10} + \sqrt{10} + 2$$

$$= 5 + 2 + \sqrt{10} + \sqrt{10}$$

$$= 7 + 2\sqrt{10}$$

3. Expand and simplify $\sqrt{5}\left(\sqrt{45} - \frac{8}{\sqrt{80}}\right)$

Solution:

$$\sqrt{5} \left(\sqrt{45} - \frac{8}{\sqrt{80}} \right) = \sqrt{5 \times 45} - \sqrt{5} \times \frac{8}{\sqrt{80}} \qquad \text{Multiply } \sqrt{5} \text{ by each term in the bracket}$$

$$= \sqrt{225} - \frac{8\sqrt{5}}{\sqrt{80}}$$

$$= 15 - \frac{8\sqrt{5}}{\sqrt{16 \times 5}}$$

$$= 15 - \frac{8\sqrt{5}}{4\sqrt{5}}$$

$$= 15 - \frac{8}{4} \qquad \text{Cancel } \sqrt{5}$$

$$= 13$$

4. Simplify:
$$\sqrt{2}\left(\sqrt{20} + \frac{4}{\sqrt{40}}\right)$$

Solution:

Note that it is easier to simplify $\sqrt{20}$ and $\sqrt{40}$ first. It is possible to solve the problem by multiplying $\sqrt{2}$ first, but this will result in larger numbers within surds.

$$\begin{split} \sqrt{2} \left(\sqrt{20} + \frac{4}{\sqrt{40}} \right) &= \sqrt{2} \left(\sqrt{4 \times 5} + \frac{4}{\sqrt{4 \times 10}} \right) & \text{Simplify } \sqrt{20} \text{ and } \sqrt{40} \\ &= \sqrt{2} \left(2\sqrt{5} + \frac{4}{2\sqrt{10}} \right) \\ &= 2\sqrt{2} \left(2\sqrt{5} + \frac{4\sqrt{2}}{2\sqrt{10}} \right) & \text{Multiply each term by } \sqrt{2} \\ &= 2\sqrt{2 \times 5} + \frac{4\sqrt{2}}{2\sqrt{10}} & \text{Simplify} \\ &= 2\sqrt{10} + \frac{2\sqrt{2}}{\sqrt{10}} \times \frac{\sqrt{10}}{\sqrt{10}} & \text{Rationalise the denominator} \\ &= 2\sqrt{10} + \frac{2\sqrt{20}}{10} \\ &= 2\sqrt{10} + \frac{\sqrt{4 \times 5}}{5} & \text{Simplify} \\ &= 2\sqrt{10} + \frac{2\sqrt{5}}{5} \\ &= 2\sqrt{10} + \frac{2\sqrt{5}}{5} \\ &= 2\sqrt{10} + \frac{2}{5}\sqrt{5} \end{split}$$

Practice

- 1. Simplify: $\sqrt{2}(\sqrt{10} + \sqrt{2})$
- 2. Expand and simplify: $\sqrt{18}(2\sqrt{5} \sqrt{3})$
- 3. Expand and simplify:

a.
$$(1 + \sqrt{2})(1 - \sqrt{2})$$

b.
$$(2 - 3\sqrt{2})(3\sqrt{2} + 1)$$

- 4. Expand and simplify: $(1 + \sqrt{2})^2$
- 5. Simplify: $\sqrt{5}\left(\sqrt{50} + \frac{5}{\sqrt{50}}\right)$

Lesson Title: Practice of surds	Theme: Numbers and Numeration
Practice Activity: PHM1-L096	Class: SSS 1

By the end of the lesson, you will be able to apply various operations to simplify expressions involving surds.

Overview

This lesson is on evaluating various expressions involving surds. The rules from previous lessons will be used to solve problems, including those for operating on and rationalising surds.

Solved Examples

- 1. Evaluate $\frac{\sqrt{75} \times \sqrt{50}}{\sqrt{80} \times \sqrt{180}}$ Solution: $\frac{\sqrt{75} \times \sqrt{50}}{\sqrt{80} \times \sqrt{180}} = \frac{\sqrt{25 \times 3} \times \sqrt{25 \times 2}}{\sqrt{16 \times 5} \times \sqrt{36 \times 5}}$ Reduce each surd to basic form $= \frac{5\sqrt{3} \times 5\sqrt{2}}{4\sqrt{5} \times 6\sqrt{5}}$ $= \frac{(5 \times 5)\sqrt{3 \times 2}}{(4 \times 6)\sqrt{5} \times \sqrt{5}}$ Multiply $= \frac{25\sqrt{6}}{24 \times 5} = \frac{5}{24}\sqrt{6}$ Simplify
- 2. If $x\sqrt{45} \sqrt{125} + 2\sqrt{5} = 0$, find *x* Solution:

$$x\sqrt{45} - \sqrt{125} + 2\sqrt{5} = 0$$

$$x\sqrt{9 \times 5} - \sqrt{25 \times 5} + 2\sqrt{5} = 0$$

$$3x\sqrt{5} - 5\sqrt{5} + 2\sqrt{5} = 0$$

$$3x\sqrt{5} - 3\sqrt{5} = 0$$

$$3x\sqrt{5} - 3\sqrt{5} = 3\sqrt{5}$$

$$x = \frac{3\sqrt{5}}{3\sqrt{5}}$$

Combine like terms

$$3x\sqrt{5} = 3\sqrt{5}$$

Transpose $-3\sqrt{5}$

$$x = 1$$

Divide throughout by $3\sqrt{5}$

3. Given $\sqrt{2} = 1.414$, evaluate $\sqrt{72} + \frac{2}{\sqrt{8}}$.

Solution:

$$\sqrt{72} + \frac{2}{\sqrt{8}} = \sqrt{36 \times 2} + \frac{2}{\sqrt{4 \times 2}}$$
 Reduce the surds to basic form
$$= 6\sqrt{2} + \frac{2}{2\sqrt{2}}$$
$$= \frac{6\sqrt{2}}{1} + \frac{1}{\sqrt{2}}$$

$$= \frac{\sqrt{2}(6\sqrt{2})+1}{\sqrt{2}}$$
Add

$$= \frac{6\times 2+1}{\sqrt{2}}$$

$$= \frac{12+1}{\sqrt{2}}$$

$$= \frac{13}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}}$$
Rationalise the denominator

$$= \frac{13\sqrt{2}}{2}$$

$$= \frac{13\times 1.414}{2}$$
Substitute $\sqrt{2} = 1.414$

$$= 9.191$$

4. Simplify $\frac{1+\sqrt{2}}{1-\sqrt{2}}$

Solution:

$$\frac{1+\sqrt{2}}{1-\sqrt{2}} = \frac{1+\sqrt{2}}{(1-\sqrt{2})} \times \frac{1+\sqrt{2}}{(1+\sqrt{2})}$$
Rationalise the denominator
$$= \frac{(1+\sqrt{2})(1+\sqrt{2})}{(1-\sqrt{2})(1-\sqrt{2})}$$
Expand and simplify the numerator
$$= \frac{1+\sqrt{2}+\sqrt{2}+(\sqrt{2})^2}{1-2}$$
Simplify
$$= \frac{1+2\sqrt{2}+2}{-1}$$

$$= \frac{3+2\sqrt{2}}{-1}$$

$$= -3 - 2\sqrt{2}$$

5. Simplify $\frac{2+\sqrt{3}}{2-\sqrt{3}} - \frac{2-\sqrt{3}}{2+\sqrt{3}}$ and write your answer in the form $a + b\sqrt{c}$. Identify *a*, *b* and *c* in your result.

Solution:

$$\frac{2+\sqrt{3}}{2-\sqrt{3}} - \frac{2-\sqrt{3}}{2+\sqrt{3}} = \frac{(2+\sqrt{3})(2+\sqrt{3})-(2-\sqrt{3})(2-\sqrt{3})}{(2-\sqrt{3})(2+\sqrt{3})}$$
Subtract the fractions
$$= \frac{[4+4\sqrt{3}+(\sqrt{3})^2]-[4-4\sqrt{3}+(\sqrt{3})^2]}{(2)^2-(\sqrt{3})^2}$$
Expand each part of the numerator
$$= \frac{(4+4\sqrt{3}+3)-(4-4\sqrt{3}+3)}{4-3}$$
Simplify
$$= \frac{7+4\sqrt{3}-7+4\sqrt{3}}{1}$$
$$= 8\sqrt{3}$$

Writing result in the form $a + b\sqrt{c}$ we have result as $0 + 8\sqrt{3}$, a = 0, b = 8 and c = 3

6. Given $\sqrt{2} = 1.414$, $\sqrt{3} = 1.732$ and $\sqrt{5} = 2.236$, evaluate: a. $\sqrt{125}(6 - \sqrt{5})$

b. $(\sqrt{8} + \sqrt{12}) - (\sqrt{18} - \sqrt{12})$ Solution: a. $\sqrt{125}(6 - \sqrt{5})$ $= \sqrt{25 \times 5}(6 - \sqrt{5})$ $= \sqrt{25} \times \sqrt{5}(6 - \sqrt{5})$ $= 5\sqrt{5}(6 - \sqrt{5})$ $= 5\sqrt{5}(6 - \sqrt{5})$ $= 5 \times 6\sqrt{5} - 5 \times \sqrt{5} \times \sqrt{5}$ $= 30\sqrt{5} - 5 \times 5$ $= 3\sqrt{5} - 25$ = 30(2.236) - 25 = 67.08 - 25= 42.08

$$\begin{array}{ll} \text{O.} & \left(\sqrt{8} + \sqrt{12}\right) - \left(\sqrt{18} - \sqrt{12}\right) \\ &= \left(\sqrt{4 \times 2} + \sqrt{4 \times 3}\right) - \left(\sqrt{9 \times 2} - \sqrt{4 \times 3}\right) \\ &= \left(2\sqrt{2} + 2\sqrt{3}\right) - \left(3\sqrt{2} - 2\sqrt{3}\right) \\ &= 2\sqrt{2} + 2\sqrt{3} - 3\sqrt{2} + 2\sqrt{3} \\ &= 2\sqrt{2} - 3\sqrt{2} + 2\sqrt{3} + 2\sqrt{3} \\ &= -\sqrt{2} + 4\sqrt{3} \\ &= -1.414 + 4(1.732) \\ &= 5.514 \end{array}$$

7. Evaluate: $\sqrt{0.48}$

Solution:

Note that when a decimal is in a surd, it can be evaluated by first converting it to a fraction.

$$\sqrt{0.48} = \sqrt{\frac{48}{100}}$$
Convert to a fraction
$$= \frac{\sqrt{48}}{\sqrt{100}}$$
Simplify
$$= \frac{\sqrt{16\times3}}{10}$$

$$= \frac{\sqrt{16}\times\sqrt{3}}{10}$$

$$= \frac{4\times\sqrt{3}}{10}$$

$$= \frac{2}{3}\sqrt{3}$$

Practice

- 1. Evaluate $\frac{\sqrt{72} \times \sqrt{48}}{\sqrt{27} \times \sqrt{128}}$ 2. If $a\sqrt{40} + \sqrt{90} - \sqrt{10} = \sqrt{1000}$, find the value of *a* 3. Given $\sqrt{5} = 2.236$, Evalute $\sqrt{125} - \frac{10}{\sqrt{500}}$ 4. Simplify $\frac{3+\sqrt{5}}{3-\sqrt{5}}$ 5. Simplify $\frac{2+\sqrt{3}}{2-\sqrt{3}} + \frac{2-\sqrt{3}}{2+\sqrt{3}}$
- 6. Evaluate $\sqrt{0.08}$

Answer Key – Term 2

Lesson Title: Powers and roots of logarithms – Numbers greater than 1 **Practice Activity:** PHM1-L049

- 1. 967.4
- 2. 132.0
- 3. 15
 4. 7.907
- 4. 7.907
- 5. 53.07
- 6. 1.841

Lesson Title: Logarithm of Numbers less than 1 Practice Activity: PHM1-L050

1. a. 4.525×10^{-1} , $\overline{1}$; b. 6.07×10^{-2} , $\overline{2}$; c. 4.98×10^{-3} , $\overline{3}$; d. 9.89×10^{-4} , $\overline{4}$ 2. a. $\overline{1}$ b. $\overline{3}$ c. $\overline{2}$ d. $\overline{5}$ 3. a. $\overline{1}.0899$ b. $\overline{2}.9943$ c. $\overline{3}.8156$ d. $\overline{4}.9538$

Lesson Title: Antilogarithm-Numbers less than 1 Practice Activity: PHM1-L051

- 1. 0.02026
- 2. 0.0001023
- 3. 0.1589
- 4. 0.0002597
- 5. 0.000001288
- 6. 0.00009997
- 7. 0.04269
- 8. 0.002160

Lesson Title: Multiplication and division of numbers less than 1-using logarithmic tables Practice Activity: PHM1-L052

- 1. 12.5
- 2. 0.0004950
- 3. 0.1129
- 4. 0.1021
- 5. 9.063
- 6. 3.109

Lesson Title: Powers and roots of logarithms – numbers less than 1 **Practice Activity:** PHM1-L053

1.	a. 0.002107	b. 1.953×10^{-9}
2.	a. 0.8548	b. 0.1044
3.	a. 0.2832	b. 0.0366
4.	a. 0.2381	b. 0.1870

Lesson Title:	Law of logarithms – Part 1
Practice Activi	ity: PHM1-L054

- 1. a. $\log_{10} 2 + \log_{10} 7$, b. $\log_{10} 3 + \log_{10} 5$, c. $\log 7 + \log 7$, d. $\log 5 + \log 5$
- 2. a. $\log_2 48$, b. $\log_5 24$, c. $\log_4 128$, d. $\log_3 36$
- 3. a. 1.4771, b. 1.7481, c. 1.8451

Lesson Title: Laws of logarithms – Part 2 Practice Activity: PHM1-L055

- 1. a. $\log_7 17 \log_7 3$; b. $\log_7 39 \log_7 5$
- 2. a. 1; b. $\log_3 \frac{16}{9}$
- 3. a. 1.18; b. -0.318; c. -0.636
- 4. a. 1.067; b. 0.7872

Lesson Title: Law of logarithms – Part 3 Practice Activity: PHM1-L056

1. a. $8 \log_3 x$; b. 3; c. 17 2. a. 1.293; b. -0.138; c. 1.795 3. a. 2; b. $\frac{2}{3}$; c. 4 4. a. 7; b. 7

Lesson Title: Define and describe sets and elements of set **Practice Activity:** PHM1-L057

- 1. a. Set of capital cities b. Set of living things c. Set of even numbers between 1 and 20,
- 2. a. {January, June, July} b. {1, 3, 5, 7, 9, 11, 13, 15, 17, 19} c. {2, 4, 6, 8},
- 3. a. Set of capital cities in West Africa b. Set of Farm animals c. Set of colours in the rainbow
- 4. a. 6 ∉ {4, 9, 16, 25.....81} b. $\frac{4}{5}$ ∉{Whole numbers} c. $\sqrt{2} \in \{\text{irrational numbers}\}$
- 5. a. n(D) = 6 b. n(X) = 6 c. n(Y) = 1

Lesson Title: Set Notation Practice Activity: PHM1-L058

- 1. A={1, 2, 3, 4, 6, 8, 12, 24}
- 2. B={10, 11, 12, 13, 14, 15}
- 3. D={ 3, 5, 7, 11, 13, 17, 19}
- 4. B={January, March, May, July, August, October, December}
- 5. $F=\{x: x \text{ is a multiple of } 10\}$
- 6. $G = \{x: x \text{ is an odd number between 0 and 10}\}$

Lesson Title: Finite and infinite set **Practice Activity:** PHM1-L059

- 1. finite; {1, 2, 3, 6, 9, 18}
- 2. infinite; {0, 1, 2, 3, 4, ...}
- 3. finite; {January, February, ..., December}
- 4. finite; {8, 16, 24, ..., 64}
- 5. infinite; {7, 14, 21, 28, ...}
- 6. finite; {12, 14, 16, ..., 38}
- 7. infinite; {51, 53, 59, ...}

Lesson Title: Null/empty, unit and universal set Practice Activity: PHM1-L060

- 1. (a) Empty set (b) Empty set (c) not an empty set (d) Empty set
- 2. (a) $D = \{May\}$ (b) $E = \{7\}$ (c) $W = \{Sunday\}$
- 3. $U = \{x: x \text{ is an integer}\}$
- 4. a. $C = \{$ Square, Rectangle, trapezium, Rhombus, kite, trapezoid $\}$
 - b. E = {Square, Rectangle, Rhombus, Parallelogram}

Lesson Title: Equivalent and equal sets Practice Activity: PHM1-L061

- 1. a. K and L are equivalent b. A and B are not equivalent c. C and T are not equivalent d. M and N are equivalent
- 2. a. V and W are not equalb. C and D are not equalc. A and B are equald. X and Y are not equale. R and S are not equalf. I and J are equal

Lesson Title: Subsets Practice Activity: PHM1-L062

- 1. $S_0 = \{a, b, c, d\}, S_1 = \{a, b, c\}, S_2 = \{b, c, d\}, S_3 = \{c, d, a\}, S_4 = \{d, a, b\}, S_5 = \{a, b\}, S_6 = \{a, c\} S_7 = \{a, d\} S_8 = \{b, c\} S_9 = \{b, d\} S_{10} = \{c, d\} S_{11} = \{a\} S_{12} = \{b\} S_{13} = \{c\} S_{14} = \{d\}, S_{15} = \Phi = \{\}$
- 2. a. W is a subset of X; b. V is a subset of X; c. T is a subset of X.
- 3. Venn diagrams:



4. $M_1 = \{2, 4, 6\}, M_2 = \{2, 4\}, M_3 = \{2, 6\}, M_4 = \{4, 6\}, M_5 = \{2\}, M_6 = \{4\}, M_7 = \{6\}, M_8 = \{ \}$

Lesson Title: Intersection of 2 sets Practice Activity: PHM1-L063

1. a. P ∩ Q {12, 24}; b. Venn diagram:



2. a. $R = \{1, 2, 3, 4, 6, 12\}, S = \{2, 3\};$ b. $R \cap S = \{2, 3\};$ c. Venn diagram:



3. a. A = {4, 8, 12, 16} B = {8, 10, 12, 14, 16, 18}; b. A \cap B = {8, 12, 16}; c. Venn diagram:



4. a. $A = \{1, 2, 3, 4, 9\}, B = \{1, 4, 5, 7, 8\}; b. A \cap B = \{1, 4\}$

Lesson Title:	Intersection of three sets
Practice Activi	ty: PHM1-L064

1. a. i{4, 8, 12}; ii. {8, 12}; iii. {8, 12, 16}; iv. {8, 12}

b. Venn diagram:



2. a. $P \cap Q = \{3, 6\}, P \cap R = \{3\}, Q \cap R = \{3\}, P \cap Q \cap R = \{3\},$

b. Venn diagram:



3. a. $M \cap L = \{ \}, M \cap T = \{3, 9, 15\}, L \cap T = \{6, 12\}, M \cap L \cap T = \{ \},$

b. Venn diagram:



4. $X \cap Y = \{1, 3, 5\}, X \cap Z = \{1, 2, 5\}, Y \cap Z = \{1, 5, 6\}, X \cap Y \cap Z = \{1, 5\}$

Lesson Title:	Disjoint sets
Practice Activity: PHM1-L065	

1. $M \cap N = \emptyset$



2. $W \cap N = \emptyset$



3. a. X={3,4,5,6}, Y={-3,-2,-1,0}; b. $X \cap Y = \emptyset$; c. Diagram:



4. $P \cap R = \emptyset$



Lesson Title: Union of two sets Practice Activity: PHM1-L066

1. $M \cup N = \{4, 5, 6, 7, 8, 10, 11\}$



2. a. i. {a, b, c, d, e,}; ii. {b, c, d, e, f}; iii. {a, b, c, d, f}; b. Venn diagrams:



- a. A∪B {Brima, Sama, Yomba, sia}; b. B∪C = {Sia, Sama}; c. A∪C = {Brima, Sama, Sia, Yomba}
- 4. a. $\{2, 3, 4, 6, 7, 8, 10\}$; b. $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} = U$; c. $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} = U$; d. $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} = U$

Lesson Title:Complement of a setPractice Activity:PHM1-L067

1. a. $U = \{4, 5, 6, 7, 8, 9, 10, 11, 12\}$ $P = \{6, 7, 8, 9\}$

b. P' = {4, 5, 10, 11, 12}

2. a. $A' = \{10, 40, 50\}$; b. $B' = \{10, 40, 60\}$; c. $A' \cap B' = \{10, 40\}$

3. a. U= {5, 10, 15, 20, 25, 30, 35, 40} $P = {5, 25, 30, 40} Q = {5, 20, 35, 40}$

b. P' = {10, 15, 20, 35}; c. Q' = {10, 15, 25, 30}; d. P' \cap Q' = {10, 15} 4.a. U = {1, 2, 3, 4, a, b, c, d, e, f, g} M = {a, 2, 3, d, e} N = {a, 1, 2, 3, b, f}

b. $M' = \{1, b, c, f, 4, g\}; c. N' = \{3, 4, c, d, e, g\}; d. M' \cap N' = \{c, g, 4\}$

Lesson Title: Real life problems involving 2 sets Practice Activity: PHM1-L068

1. a. Venn diagram:



- b. 30 pupils c. 40 played Lawn tennis only
- 2. a. Venn diagram:



b. i. 50 ii. 120 iii. 110

3. a. Venn diagram:



b. i. 19 ii. 8 iii. 13 4. a. 30 b. i. 60 ii. 10

Lesson Title:	Real life problems involving 3 sets – Part 1
Practice Activ	ity: PHM1-L069

- 1. a. 33; b. 12
- 2. a. Venn diagram:



b. i. 1 ii. 3

Lesson Title: Real life problems involving 3 sets – Part 2 Practice Activity: PHM1-L070

1. a. Venn diagram:



b. i. 8; ii. 24; iii. 20







3. a. Venn diagram:



b. i. 4; ii. 6
Lesson Title: Use of variables Practice Activity: PHM1-L071

- 1. a. *r* b. *m*, *n* c. *x u*, *t* and *a*
- 2. a. coefficients are -5 and 6; constant is -10
 - b. coefficients are $\frac{4}{5}$ and -1; constant is 7
 - c. coefficients are 1 and -1; constant is 10
- 3. a. 60x; b. 100y, c. y + 2x or 2x + y,

4. a. *x* = 13; b. *a* = 4; c. *b* = 3, d. *z* = 6

Lesson Title: Simplification – grouping terms Practice Activity: PHM1-L072

- 1. 2x + y
- 2. -2a + 7b
- 3. m + 14n
- 4. 5xy xp
- 5. 12mn m
- 6. 2pq + 8qr
- 7. $8p^2q 3pq^2$
- 8. 9xyz xz
- 9. 2ab + 4a + 4

Lesson Title: Simplification – removing brackets Practice Activity: PHM1-L073

- 1. 5x 20
- 2. -21y + 28
- 3. 7m 2n
- 4. 2u 6uv
- 5. -3x 3y
- 6. -26m + 14n

Lesson Title: Simplification – expanding brackets Practice Activity: PHM1-L074

- 1. $7x^2 + 30x + 8$
- 2. $p^2 11p + 30$

- 3. $m^2 + m 6$ 4. $x^2 - 8x + 16$ 5. $n^2 + 6n + 9$ 6. 8mn - 2n + 19m - 27. $3x^3 - 12x^2 + 12x$
- 8. $4x^2 + 48x + 144$

Lesson Title: Factoring – common factors Practice Activity: PHM1-L075

1. 9(ab-3)2. x(x-4)3. $a(a^2+b)$ 4. $4x^2(4x^2-2x+1)$ 5. $9a^2(9a+1)$ 6. $12x^2(x^2-3x+4)$ 7. $3s^2t^2(6s^3t^4-2s^2t+1)$ 8. $5uv^2(v-5+3u)$

Lesson Title: Factoring – grouping	
Practice Activity: PHM1-L076	

1.
$$(x-1)(a+1)$$
,
2. $2(3g-h)(2e-f)$,
3. $(d-e)(c-d)$,
4. $(1+5a)(p+q)$,
5. $(x+2)(x-5)$,
6. $(2a-1)(3a+2)$,
7. $(x+2)(x+3)$,
8. $(3-2d)(x+y)$

Lesson Title: Substitution of values	
Practice Activity: PHM1-L077	

1. -12, 2. a. $\frac{4}{25}$, b. 231, c. $12\frac{1}{4}$, 3. $3\frac{9}{19}$, 4. a. 15, b. 35 Lesson Title: Addition of algebraic fractions Practice Activity: PHM1-L078

1. a.
$$a$$
 b. $\frac{7m}{9}$, c. $\frac{8x}{7}$, d. $\frac{x+2}{2}$
2. a. $\frac{8a}{15}$, b. $\frac{5x}{4}$, c. $\frac{11n}{10}$, d. $\frac{3x}{8}$
3. a. $\frac{13a-1}{12}$, b. $\frac{29n+8}{28}$, c. $\frac{3x-1}{6}$, d. $\frac{3x+2}{10}$

Lesson Title: Subtractions of algebraic fractions	
Practice Activity: PHM1-L079	

1.	a. $\frac{x}{4}$,	b. $\frac{2y-3}{3}$,	C. $\frac{2x+1}{5}$,	d. – <i>a</i>
2.	a. $\frac{a}{12}$,	b. $-\frac{x}{5}$,	C. $\frac{11x}{6}$,	d. $-\frac{x}{20}$
3.	a. $\frac{13y-6}{6}$,	b. $\frac{a-6}{6}$,	C. $\frac{5y+11}{12}$,	d. $\frac{8y+5}{12}$

Lesson Title: Linear equations	
Practice Activity: PHM1-L080	

1.
$$m = -3$$

2. $y = -\frac{1}{2}$
3. $n = 20$
4. $y = 1\frac{2}{3}$
5. $x = 1$
6. $y = 1\frac{1}{4}$
7. $x = -1$
8. $m = 0$
9. $y = -5\frac{1}{2}$

Lesson Title: Linear equations with brackets Practice Activity: PHM1-L081

1. a = 52. $m = \frac{1}{2}$ 3. x = -104. y = -1 5. $a = -\frac{1}{4}$ 6. x = 27. $p = 2\frac{1}{2}$

Lesson Title: Linear equations with fractions	
Practice Activity: PHM1-L082	

1. x = 152. a = 203. $y = 4\frac{4}{5}$ 4. $x = 1\frac{1}{9}$ 5. x = 186. k = 337. $p = 4\frac{1}{2}$

Lesson Title:	Word problems	
Practice Activi	ty: PHM1-L083	

1. a.
$$\frac{15+x}{4} = 80 - x$$
; b. 61

- 2. a. $\frac{x}{20}$; b. 160
- 3. 6 years, 17 years
- 4. Le 3,200.00, Le 4,150.00
- 5. 11 mangoes, 48 mangoes

Lesson Title: Substitution in formulae Practice Activity: PHM1-L084

- 1. a. 6; b. 2; c. 31; d. 8
- 2. 5.56 or $5\frac{5}{9}$
- 3. 15
- 4. 22cm
- 5. 36,000
- 6. 2,355 cm²

Lesson Title: Change of subject- Part 1 Practice Activity: PHM1-L085

1. a.
$$b = \frac{a}{13}$$
; b. $b = a + 3$; c. $b = 4a$; d. $b = \frac{a}{2} + 4$
2. $a = \frac{p}{2} - b$,
3. a. $y = \frac{rt - 4d}{5c - q}$, b. $y = \frac{r + s}{3p - a}$, c. $y = \frac{-9x}{k - 4}$
4. a. $u = \frac{1 + 2v}{v - 1}$, b. $u = \frac{w}{mh - 3}$, c. $u = \frac{nv - 2v}{2k}$
5. $F = \frac{9}{5}c + 32$

Lesson Title:	Change of Subject – Part 2
Practice Activ	ity: PHM1-L086

1.
$$x = 144$$

2. $r = \sqrt{p-1}$
3. $x = 9y^2 - 1; x = 35$ when $y = -2$.
4. $u = \frac{mv^2}{F}$
5. $t = \pm \sqrt{\frac{2x-2uf}{g}}$
6. $r = \pm \sqrt{\frac{3v}{\pi}}$
7. $u = \frac{c^2b-a}{c^2-1}$

Lesson Title: Reduction to Base Forms of Surds Practice Activity: PHM1-L087

- 1. a. $4\sqrt{2}$; b. $2\sqrt{10}$; c. $5\sqrt{2}$; d. $3\sqrt{3}$
- 2. a. $12\sqrt{3}$; b. $6\sqrt{5}$; c. $25\sqrt{2}$; d. $4\sqrt{15}$
- 3. a. $25\sqrt{10}$; b. $9\sqrt{3}$; c. $30\sqrt{10}$; d. $20\sqrt{2}$

Lesson Title: Addition and Subtraction of Surds - Part 1 Practice Activity: PHM1-L088

- 1. $4\sqrt{2}$
- 2. $7\sqrt{5}$
- 3. $2\sqrt{3}$

4. 0 5. $3\sqrt{2}$ 6. $\sqrt{3}$

7. $11\sqrt{2}$

Lesson Title: Addition and Subtraction of Surds – Part 2 Practice Activity: PHM1-L089

1. $\frac{5}{2}\sqrt{7}$ 2. $3\sqrt{2}$ 3. $11\sqrt{2}$ 4. $8\sqrt{3} + \sqrt{6}$ 5. $9\sqrt{5}$ 6. $4\sqrt{7}$ 7. n = 1

Lesson Title: Properties of Surds	
Practice Activity: PHM1-L090	

1. a. 4√2;	b. 4√3;	c. 2√6
2. a. 1 ¹ / ₂ ;	b. $\frac{5\sqrt{2}}{3};$	c. $1\frac{1}{4}$
3. a. − <i>a</i> √6;	b. $\sqrt{2};$	c. $\sqrt{3}$
4. a. 8√2;	b. 8√ 6 ;	c. $a^2 \sqrt{a}$ (d) x^2

Lesson Title:	Multiplication of Surds – Part 1
Practice Activ	ity: PHM1-L091

1.	a. 105;	b. 16√10
2.	a. 32;	b. $144\sqrt{6}$
3.	a. 32√ 3 ;	b. $50\sqrt{15}$
4.	a. 18;	b. 48√3

Lesson Title: Multiplication of Surds – Part 2 Practice Activity: PHM1-L092

1. $10\sqrt{5}$ 2. $3\sqrt{2} - 4$ 3. $9\sqrt{6} - 6$ 4. $12\sqrt{5} - 30$ 5. $24 + 12\sqrt{2}$ 6. $1\frac{1}{5}$ 7. $31 + 12\sqrt{3}$

Lesson Title: Rationalisation of the Denominator of Surds – Part 1 Practice Activity: PHM1-L093

1. $\frac{3\sqrt{2}}{4}$ 2. $\frac{\sqrt{35}}{7}$ 3. $\sqrt{2}$ 4. $\frac{\sqrt{6}}{4}$ 5. $\frac{\sqrt{6}}{3}$ 6. $\frac{\sqrt{10}}{3}$ 7. $2\sqrt{6}$ 8. $\sqrt{7}$ 9. $\sqrt{2}$ 10. $\frac{2\sqrt{6}}{3}$

Lesson Title: Rationalisation of the Denominator of Surds – Part 2 **Practice Activity:** PHM1-L094

1. a.
$$\frac{12+3\sqrt{3}}{13}$$
; b. $\frac{3-\sqrt{2}}{7}$; c. $\frac{25-5\sqrt{2}}{23}$; d. $\frac{42+12\sqrt{2}}{41}$; e. $\frac{14}{13} - \frac{7}{26}\sqrt{3}$,
2. a. $\frac{3}{4}\sqrt{2} - \frac{1}{2}\sqrt{6}$; b. $\frac{5}{3} + \frac{\sqrt{15}}{9}$; c. $\frac{4}{5}\sqrt{5} + \frac{\sqrt{15}}{5}$,
3. $\frac{1}{4}\sqrt{2} + \frac{1}{6}\sqrt{3}$

Lesson Title: Expansion and Simplification of Surds Practice Activity: PHM1-L095

1. $2 + 2\sqrt{5}$ 2. $6\sqrt{10} - 3\sqrt{6}$ 3. a. -1; b. $-16 + 3\sqrt{2}$ 4. $3 + 2\sqrt{2}$ 5. $\frac{11}{2}\sqrt{10}$

Lesson Title:	Practice of Surds
Practice Activi	ity: PHM1-L096

- 1. 1 2. 4
- 3. 10.7328

- 4. $\frac{7}{2} + \frac{3}{2}\sqrt{5}$ 5. 14 6. $\frac{1}{5}\sqrt{2} \text{ or } \frac{\sqrt{2}}{5}$

Appendix I: Logarithm Table

_	0	1	2	3	4	5	6	7	0	0	1	-			Die				1
	v		-	3	-	2	0	· ·	•	9	1	2	3	A	Dille	6	7	8	0
10	0000	0043	0086	0128	0170	0212	0253	0294	0334	0374	4	8	12	17	21	25	20	32	27
11	0414	0453	0492	0531	0569	0607	0645	0682	0719	0755	4	8	11	15	19	23	26	30	34
12	0792	0828	0864	0899	0934	0969	1004	1038	1072	1106	3	7	10	14	17	21	24	28	31
14	1139	1492	1206	1239	12/1	1303	1335	1367	1399	1430	3	6	10	13	16	19	23	26	29
15	1761	1790	1818	1847	1875	1903	1931	1959	1987	2014	3	6	8	11	14	17	20	22	25
16	2041	2068	2095	2122	2148	2175	2201	2227	2253	2279	3	5	8	11	13	16	18	21	24
17	2304	2330	2355	2380	2405	2430	2455	2480	2504	2529	2	5	7	10	12	15	17	20	22
19	2788	2810	2833	2856	2878	2900	2923	2945	2967	2989	2	4	7	9	11	13	16	18	21
20	3010	3032	3054	3075	3096	3118	3139	3160	3181	3201	2	4	6	8	11	13	15	17	19
21	3222	3243	3263	3284	3304	3324	3345	3365	3385	3404	2	4	6	8	10	12	14	16	18
23	3617	3636	3655	3674	3692	3711	3729	3747	3766	3784	2	4	6	8	9	12	14	15	17
24	3802	3820	3838	3856	3874	3892	3909	3927	3945	3962	2	4	5	7	9	11	12	14	16
25	3979	3997	4014	4031	4048	4065	4082	4099	4116	4133	2	3	5	7	9	10	12	14	15
27	4150	4100	4185	4200	4210	4232	4249	4265	4281	4298	2	3	5	6	8	10	11	13	15
28	4472	4487	4502	4518	4533	4548	4564	4579	4594	4609	2	3	5	6	8	9	11	12	14
29	4624	4639	4654	4669	4683	4698	4713	4728	4742	4757	1	3	4	6	7	9	10	12	13
30	4771	4786	4800	4814	4829	4843	4857	4871	4886	4900	1	3	4	6	7	9	10	11	13
32	5051	5065	5079	5092	5105	5119	5132	5145	5159	5038	1	3	4	5	7	8	10	11	12
33	5185	5198	5211	5224	5237	5250	5263	5276	5289	5302	i	3	4	5	6	8	9	10	12
34	5315	5328	5340	5353	5366	5378	5391	5403	5416	5428	1	3	4	5	6	8	9	10	11
35	5441	5453	5465	5478	5490	5502	5514	5527	5539	5551	1	2	4	5	6	7	9	10	11
37	5682	5694	5705	5717	5729	5740	5752	5763	5775	5786	L i	2	4	5	6	7	8	9	10
38	5798	5809	5821	5832	5843	5855	5866	5877	5888	5899	1	2	3	5	6	7	8	9	10
39	5911	5922	5933	5944	5955	5966	5977	5988	5999	6010	1	2	3	4	5	7	8	9	10
40	6021	6031	6042	6053	6064	6075	6085	6096	6107	6117	1	2	3	4	5	6	8	9	10
41	6232	6243	6253	6263	6274	6284	6294	6304	6314	6325	1	2	3	4	5	6	7	8	9
43	6335	6345	6355	6365	6375	6385	6395	6405	6415	6425	i	2	3	4	5	6	7	8	9
44	6435	6444	6454	6464	6474	6484	6493	6503	6513	6522	1	2	3	4	5	6	7	8	9
45	6532	6542	6551	6561	6571	6580	6590	6599	6609	6618	1	2	3	4	5	6	7	8	9
40	6721	6730	6739	6749	6758	6767	6776	6785	6794	6803	1	2	3	4	5	5	6	7	8
48	6812	6821	6830	6839	6848	6857	6866	6875	6884	6893	i	2	3	4	4	5	6	7	8
49	6902	6911	6920	6928	6937	6946	6955	6964	6972	6981	1	2	3	4	4	5	6	7	8
50	6990	6998	7007	7016	7024	7033	7042	7050	7059	7067	1	2	3	3	4	5	6	7	8
52	7160	7168	7177	7185	7193	7202	7210	7218	7226	7235	1	2	2	3	4	5	6	7	7
53	7243	7251	7259	7267	7275	7284	7292	7300	7308	7316	1	2	2	3	4	5	6	6	7
54	7324	7332	7340	7348	7356	7364	7372	7380	7388	7396	1	2	2	3	4	5	6	6	7
55	7404	7412	7419	7427	7435	7443	7451	7459	7466	7474	1	2	2	3	4	5	5	6	7
57	7559	7566	7574	7582	7515	7597	7604	7612	7619	7627		2	2	3	4	5	5	6	7
58	7634	7642	7649	7657	7664	7672	7679	7686	7694	7701	1	1	2	3	4	4	5	6	7
59	7709	7716	7723	7731	7738	7745	7752	7760	7767	7774	1	1	2	3	4	4	5	6	7
60	7782	7789	7796	7803	7810	7818	7825	7832	7839	7846	1	1	2	3	4	4	5	6	6
62	7924	7931	7938	7945	7952	7959	7966	7903	7910	7987	1	1	2	3	4	4	5	6	6
63	7993	8000	8007	8014	8021	8028	8035	8041	8048	8055	i	1	2	3	3	4	5	5	6
64	8062	8069	8075	8082	8089	8096	8102	8109	8116	8122	1	1	2	3	3	4	5	5	6
65	8129	8136	8142	8149	8156	8162	8169	8176	8182	8189		1	2	3	3	4	5	5	6
67	8261	8267	8274	8280	8287	8293	8299	8306	8312	8319	1	1	2	3	3	4	5	5	6
68	8325	8331	8338	8344	8351	8357	8363	8370	8376	8382	1	1	2	3	3	4	4	5	6
69	8388	8395	8401	8407	8414	8420	8426	8432	8439	8445	1	1	2	2	3	4	4	5	6
71	8513	8457	8463	8470	84/6	8482	8488	8494	8500	8500	1	1	2	2	3	4	4	5	5
72	8573	8579	8585	8591	8597	8603	8609	8615	8621	8627	î	î	2	2	3	4	4	5	5
73	8633	8639	8645	8651	8657	8663	8669	8675	8681	8686	1	1	2	2	3	4	4	5	5
74	8692	8698	8704	8710	8716	8722	8727	8733	8739	8745		1	2	2	3	4	4	5	5
76	8808	8814	8820	8825	8831	8837	8842	8848	8854	8859	1	1	2	2	3	3	4	5	5
77	8865	8871	8876	8882	8887	8893	8899	8904	8910	8915	1	1	2	2	3	3	4	4	5
78	8921	8927	8932	8938	8943	8949	8954	8960	8965	8971	1	1	2	2	3	3	4	4	5
80	8976	8982	8987	8993	8998	9004	9009	9015	9020	9025	1	1	2	2	3	3	4	4	5
81	9085	9090	9096	9101	9106	9112	9117	9122	9128	9133	1	1	2	2	3	3	4	4	5
82	9138	9143	9149	9154	9159	9165	9170	9175	9180	9186	1	1	2	2	3	3	4	4	5
83	9191	9196	9201	9206	9212	9217	9222	9227	9232	9238	1	1	2	2	3	3	4	4	5
84	9243	9248	9253	9258	9263	9269	9274	9279	9284	9289	1	1	2	2	3	3	4	4	5
85	9294	9299	9304	9309	9315	9320	9325	9330	9335	9340	1	1	2	2	3	3	4	4	5
87	9395	9400	9405	9410	9415	9420	9425	9430	9435	9440	Ó	î	î	2	2	3	3	4	4
88	9445	9450	9455	9460	9465	9469	9474	9479	9484	9489	0	1	1	2	2	3	3	4	4
89	9494	9499	9504	9509	9513	9518	9523	9528	9533	9538	0	1	1	2	2	3	3	4	4
90	9542	9547	9552	9557	9562	9566	9571	9576	9581	9586	0	1	1	2	2	3	3	4	4
92	9638	9643	9647	9652	9657	9661	9666	9671	9675	9680	0	1	i	2	2	3	3	4	4
93	9685	9689	9694	9699	9703	9708	9713	9717	9722	9727	0	1	1	2	2	3	3	4	4
94	9731	9736	9741	9745	9750	9754	9759	9763	9768	9773	0	1	1	2	2	3	3	4	4
95	9823	9827	9/80	9836	9795	9800	9805	9809	9814	9818	0	1	1	2	2	3	3	4	4
97	9868	9872	9877	9881	9886	9890	9894	9899	9903	9908	0	i	î	2	2	3	3	4	4
98	9912	9917	9921	9926	9930	9934	9939	9943	9948	9952	0	1	1	2	2	3	3	4	4
99	9956	9961	9965	9969	9974	9978	9983	9987	9991	9996	0	1	1	2	2	3	3	3	4

COMMON LOGARITHM TABLE

Appendix II: Anti-Logarithm Table

FRITHLOGARTAIM

ANTI-LOGARITHMS FRITHLOGARTAIM

ANTI-LOGARITHMS

	0	1	2	3	4	5	6	7	8	9	1 2	3	4	5	6	7	8	9	Γ	0	1	
-00	1000	1002	1005	1007	1009	1012	1014	1016	1019	1021	0.0	1	1	1	1	2	2	2	-50	3162	3170	31
-01 -02 -03	1023 1047 1072	1026 1050 1074	1028 1052 1076	1030 1054 1079	1033 1057 1081	1035 1059 1084	1038 1062 1086	1040 1064 1089	1042 1067 1091	1045 1069 1094	000000		1 1 1	111	1 1 1	222	222	2222	·51 ·52 ·53	3236 3311 3388	3243 3319 3396	32 33 34
-04 -05 -06	1096 1122 1148	1099 1125 1151	1102 1127 1153	1104 1130 1156	1107 1132 1159	1109 1135 1161	1112 1138 1164	1114 1140 1167	1117 1143 1169	1119 1146 1172	01	1 1 1	1 1 1	1 1 1	2 2 1	222	222	2222	-54 -55 -56	3467 3548 3631	3475 3556 3639	34 35 36
-07 -08 -09	1175 1202 1230	1178 1205 1233	1180 1208 1236	1183 1211 1239	1186 1213 1242	1189 1216 1245	1191 1219 1247	1194 1222 1250	1197 1225 1253	1199 1227 1256	01	1	1 1 1	1 1 1	222	222	222	233	-57 -58 -59	3715 3802 3890	3724 3811 3899	37 38 39
-10	1259	1262	1265	1268	1271	1274	1276	1279	1282	1285	0 1	1	1	1	2	2	2	3	-60	3981	3990	39
·11 ·12 ·13	1288 1318 1349	1291 1321 1352	1294 1324 1355	1297 1327 1358	1300 1330 1361	1303 1334 1365	1306 1337 1368	1309 1340 1371	1312 1343 1374	1315 1346 1377	0101	1 1 1	1 1 1	222	222	222	223	300	-61 -62 -63	4074 4169 4266	4083 4178 4276	40 41 42
-14 -15 -16	1380 1413 1445	1384 1416 1449	1387 1419 1452	1390 1422 1455	1393 1426 1459	1396 1429 1462	1400 1432 1466	1403 1435 1469	1406 1439 1472	1409 1442 1476	010101	1 1 1	1 1 1	222	222	222	333	333	-64 -65 -66	4365 4467 4571	4375 4477 4581	43 44 45
-17 -18 -19	1479 1514 1549	1483 1517 1552	1486 1521 1556	1489 1524 1560	1493 1528 1563	1496 1531 1567	1500 1535 1570	1503 1538 1574	1507 1542 1578	1510 1545 1581	0101	1 1 1	1 1 1	222	222	223	300	333	-67 -68 -69	4677 4786 4898	4688 4797 4909	46 48 49
-20	1585	1589	1592	1596	1600	1603	1607	1611	1614	1618	01	1	1	2	2	3	3	3	•70	5012	5023	50
·21 ·22 ·23	1622 1660 1698	1626 1663 1702	1629 1667 1706	1633 1671 1710	1637 1675 1714	1641 1679 1718	1644 1683 1722	1648 1687 1726	1652 1690 1730	1656 1694 1734	010101	1 1 1	222	222	222	333	333	334	·71 ·72 ·73	5129 5248 5370	5140 5260 5383	51 52 53
·24 ·25 ·26	1738 1778 1820	1742 1782 1824	1746 1786 1828	1750 1791 1832	1754 1795 1837	1758 1799 1841	1762 1803 1845	1766 1807 1849	1770 1811 1854	1774 1816 1858	0101	1 1 1	222	222	223	333	333	444	-74 -75 -76	5495 5623 5754	5508 5636 5768	55 56 57
·27 ·28 ·29	1862 1905 1950	1866 1910 1954	1871 1914 1959	1875 1919 1963	1879 1923 1968	1884 1928 1972	1888 1932 1977	1892 1936 1982	1897 1941 1986	1901 1945 1991	0101	1 1 1	222	222	333	333	344	444	·77 ·78 ·79	5888 6026 6166	5902 6039 6180	59 60 61
-30	1995	2000	2004	2009	2014	2018	2023	2028	2032	2037	0 1	1	2	2	3	3	4	4	-80	6310	6324	63
·31 ·32 ·33	2042 2089 2138	2046 2094 2143	2051 2099 2148	2056 2104 2153	2061 2109 2158	2065 2113 2163	2070 2118 2168	2075 2123 2173	2080 2128 2178	2084 2133 2183	01 01 01	1 1	2222	222	3333	333	444	444	-81 -82 -83	6457 6607 6761	6471 6622 6776	64 66 67
·34 ·35 ·36	2188 2239 2291	2193 2244 2296	2198 2249 2301	2203 2254 2307	2208 2259 2312	2213 2265 2317	2218 2270 2323	2223 2275 2328	2228 2280 2333	2234 2286 2339	1 1 1 1 1 1	222	222	333	333	444	444	555	-84 -85 -86	6918 7079 7244	6934 7096 7261	69 71 72
-37 -38 -39	2344 2399 2455	2350 2404 2460	2355 2410 2466	2360 2415 2472	2366 2421 2477	2371 2427 2483	2377 2432 2489	2382 2438 2495	2388 2443 2500	2393 2449 2506	11 11 11	222	222	333	200	444	445	5555	-87 -88 -89	7413 7586 7762	7430 7603 7780	74
-40	2512	2518	2523	2529	2535	2541	2547	2553	2559	2564	11	2	2	3	4	4	5	5	·90	7943	7962	79
·41 ·42 ·43	2570 2630 2692	2576 2636 2698	2582 2642 2704	2588 2649 2710	2594 2655 2716	2600 2661 2723	2606 2667 2729	2612 2673 2735	2618 2679 2742	2624 2685 2748	11 11 11	222	223	333	444	444	555	566	·91 ·92 ·93	8128 8318 8511	8147 8337 8531	81 83 85
-44 -45 -46	2754 2818 2884	2761 2825 2891	2767 2831 2897	2773 2838 2904	2780 2844 2911	2786 2851 2917	2793 2858 2924	2799 2864 2931	2805 2871 2938	2812 2877 2944	111111	222	333	333	444	455	555	666	-94 -95 -96	8710 8913 9120	8730 8933 9141	87 89 91
-47 -48 -49	2951 3020 3050	2958 3027 3097	2965 3034 3105	2972 3041 3112	2970 3048 3119	2985 3055 3126	2992 3062 3133	2999 3069 3141	3006 3076 3148	3013 3083 3155	11 11 11	222	333	344	444	555	566	6	-97 -98 -99	9333 9550 9772	9354 9572 9795	93 95 98

	Source and the second																		
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
-50	3162	3170	3177	3184	3192	3199	3206	3214	3221	3228	1	1	2	3	4	4	5	6	1
-51 -52 -53	3236 3311 3388	3243 3319 3396	3251 3327 3404	3258 3334 3412	3266 3342 3420	3273 3350 3428	3281 3357 3436	3289 3365 3443	3296 3373 3451	3304 3381 3459	1 1 1	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	2 2 2	333	444	5 5 5	5 5 6	666	777
-54 -55 -56	3467 3548 3631	3475 3556 3639	3483 3565 3648	3491 3573 3656	3499 3581 3664	3508 3589 3673	3516 3597 3681	3524 3606 3690	3532 3614 3698	3540 3622 3707	1 1 1	2 2 2	2	333	444	555	666	677	778
-57 -58 -59	3715 3802 3890	3724 3811 3899	3733 3819 3908	3741 3828 3917	3750 3837 3926	3758 3846 3936	3767 3855 3945	3776 3864 3954	3784 3873 3963	3793 3882 3972	1 1 1	2 2 2	3	344	4 4 5	555	666	7777	00 00 00
-60	3981	3990	3999	4009	4018	4027	4036	4046	4055	4064	1	2 :	3	4	5	6	6	7	8
61 62 63	4074 4169 4266	4083 4178 4276	4093 4188 4285	4102 4198 4295	4111 4207 4305	4121 4217 4315	4130 4227 4325	4140 4236 4335	4150 4246 4345	4159 4256 4355	1 1 1	2 2 2	3	444	555	666	7 7 7	888	999
-64 -65 -66	4365 4467 4571	4375 4477 4581	4385 4487 4592	4395 4498 4603	4406 4508 4613	4416 4519 4624	4426 4529 4634	4436 4539 4645	4446 4550 4656	4457 4560 4667	1 1 1	2 :	3	444	555	666	7 7 7	889	9 9 10
-67 -68 -69	4677 4786 4898	4688 4797 4909	4699 4808 4920	4710 4819 4932	4721 4831 4943	4732 4842 4955	4742 4853 4966	4753 4864 4977	4764 4875 4989	4775 4887 5000	1 1 1	2 2 2	333	445	566	7 7 7	8 8 8	999	10 10 10
70	5012	5023	5035	5047	5058	5070	5082	5093	5105	5117	1	2	4	5	6	7	8	9	11
71	5129 5248 5370	5140 5260 5383	5152 5272 5395	5164 5284 5408	5176 5297 5420	5188 5309 5433	5200 5321 5445	5212 5333 5458	5224 5346 5470	5236 5358 5483	1 1 1	2 .	4	555	666	7 7 8	8 9 9	10 10 10	11 11 11
-74 -75 -76	5495 5623 5754	5508 5636 5768	5521 5649 5781	5534 5662 5794	5546 5675 5808	5559 5689 5821	5572 5702 5834	5585 5715 5848	5598 5728 5861	5610 5741 5875	1 1 1	333	4	555	677	888	9 9 9	10 10 11	12 12 12
-77 -78 -79	5888 6026 6166	5902 6039 6180	5916 6053 6194	5929 6067 6209	5943 6081 6223	5957 6095 6237	5970 6109 6252	5984 6124 6266	5998 6138 6281	6012 6152 6295	1 1 1	3 .	444	566	7777	889	10 10 10	11 11 11	12 13 13
80	6310	6324	6339	6353	6368	6383	6397	6412	6427	6442	1	3.	•	6	1	9	10	12	13
-81 -82 -83	6457 6607 6761	6471 6622 6776	6486 6637 6792	6501 6653 6808	6516 6668 6823	6531 6683 6839	6546 6699 6855	6561 6714 6871	6577 6730 6887	6592 6745 6902	222	333	5	666	8 8 8	999	11 11 11	12 12 13	14 14 14
-84 -85 -86	6918 7079 7244	6934 7096 7261	6950 7112 7278	6966 7129 7295	6982 7145 7311	6998 7161 7328	7015 7178 7345	7031 7194 7362	7047 7211 7379	7063 7228 7396	222	3 3 3	5	6 7 7	888	10 10 10	11 12 12	13 13 13	15 15 15
-87 -88 -89	7413 7586 7762	7430 7603 7780	7447 7621 7798	7464 7638 7816	7482 7656 7834	7499 7674 7852	7516 7691 7870	7534 7709 7889	7551 7727 7907	7568 7745 7925	2 2 2	3 4 4	5	7 7 7	9 9 9	10 11 11	12 12 13	14 14 14	16 16 16
-90	7943	7962	7980	7998	8017	8035	8054	8072	8091	8110	2	4 (5	7	9	11	13	15	17
-91 -92 -93	8128 8318 8511	8147 8337 8531	8166 8356 8551	8185 8375 8570	8204 8395 8590	8222 8414 8610	8241 8433 8630	8260 8453 8650	8279 8472 8670	8299 8492 8690	2222	4 6 4 6	5	888	9 10 10	11 12 12	13 14 14	15 15 16	17 17 18
-94 -95 -96	8710 8913 9120	8730 8933 9141	8750 8954 9162	8770 8974 9183	8790 8995 9204	8810 9016 9226	8831 9036 9247	8851 9057 9268	8872 9078 9290	8892 9099 9311	2 2 2	4 (4 (5	888	10 10 11	12 12 13	14 15 15	16 17 17	18 19 19
-97 -98 -99	9333 9550 9772	9354 9572 9795	9376 9594 9817	9397 9616 9840	9419 9638 9863	9441 9661 9886	9462 9683 9908	9484 9705 9931	9506 9727 9954	9528 9750 9977	2 2 2	4 .	7	9 9 9	11 11 11	13 13 14	15 16 16	17 18 18	20 20 20





IN PARTNERSHIP WITH





NOT FOR SALE

Document information:

Leh Wi Learn (2018). "Maths, SeniorSecondarySchool Year 1, Term 2, pupil handbook." A resource produced by the Sierra Leone Secondary Education Improvement Programme (SSEIP). DOI: 10.5281/zenodo.3745342.

Document available under Creative Commons Attribution 4.0, https://creativecommons.org/licenses/by/4.0/.

Uploaded by the EdTech Hub, https://edtechhub.org. For more information, see https://edtechhub.org/oer. Archived on Zenodo: April 2020. DOI: 10.5281/zenodo.3745342

Please attribute this document as follows:

Leh Wi Learn (2018). "Maths, SeniorSecondarySchool Year 1, Term 2, pupil handbook." A resource produced by the Sierra Leone Secondary Education Improvement Programme (SSEIP). DOI 10.5281/zenodo.3745342. Available under Creative Commons Attribution 4.0 (https://creativecommons.org/licenses/by/4.0/). A Global Public Good hosted by the EdTech Hub, https://edtechhub.org. For more information, see https://edtechhub.org/oer.